

X-16618  
BA

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
15 March 2001 (15.03.2001)

PCT

(10) International Publication Number  
**WO 01/18172 A2**

- (51) International Patent Classification<sup>7</sup>: C12N Oaks, CA 91360 (US). LIU, Benxian [CN/US]; 2528 Rutland Place, Thousand Oaks, CA 91362 (US).
- (21) International Application Number: PCT/US00/24373
- (22) International Filing Date: 5 September 2000 (05.09.2000) (74) Agent: NOONAN, Kevin, E.; McDonnell Boehnen Hulbert & Berghoff, 300 South Wacker Drive, Suite 3200, Chicago, IL 60606 (US).
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
09/391,861 7 September 1999 (07.09.1999) US  
09/644,052 23 August 2000 (23.08.2000) US
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:  
US 09/391,861 (CON)  
Filed on 7 September 1999 (07.09.1999)  
US 09/644,052 (CON)  
Filed on 23 August 2000 (23.08.2000)
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): AMGEN, INC. [US/US]; One Amgen Center Drive, Thousand Oaks, CA 91320 (US).
- Published:  
— Without international search report and to be republished upon receipt of that report.
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): THOMASON, Arlen, Read [US/US]; 2298 Watertown Court, Thousand Oaks, CA 91320 (US).
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/18172 A2

(54) Title: FIBROBLAST GROWTH FACTOR-LIKE POLYPEPTIDES

(57) Abstract: The present invention provides novel Fibroblast Growth Factor-like (FGF-like) polypeptides and nucleic acid molecules encoding the same. The invention also provides vectors, host cells, antibodies and methods for producing FGF-like polypeptides. Also provided for are methods for the diagnosis and treatment of diseases associated with FGF-like polypeptides.

- 1 -

## FIBROBLAST GROWTH FACTOR-LIKE POLYPEPTIDES

### Field of the Invention

The present invention relates to novel Fibroblast Growth Factor-like (FGF-like) polypeptides and nucleic acid molecules encoding the same. The invention also relates to vectors, host cells, antibodies and methods for producing FGF-like polypeptides. Also provided for are methods for the diagnosis and treatment of diseases associated with FGF-like polypeptides.

### 10 Background of the Invention

Technical advances in the identification, cloning, expression, and manipulation of nucleic acid molecules have greatly accelerated the discovery of novel therapeutics based upon deciphering the human genome. Rapid nucleic acid sequencing techniques can now generate sequence information at unprecedented rates and, coupled with computational analyses, allow the assembly of overlapping sequences into entire genomes and the identification of polypeptide-encoding regions. Comparison of a predicted amino acid sequence against a database compilation of known amino acid sequences can allow one to determine the extent of homology to previously identified sequences and/or structural landmarks. Cloning and expression of a polypeptide-encoding region of a nucleic acid molecule provides a polypeptide product for structural and functional analysis. Manipulation of nucleic acid molecule and encoded polypeptides to give variants and derivatives thereof may confer advantageous properties on a product for use as a therapeutic.

25 In spite of the significant technical advances in genome research over the past decade, the potential for development of novel therapeutics based on the human genome is still largely unrealized. Genes encoding potentially beneficial protein therapeutics, or those encoding polypeptides that may act as "targets" for therapeutic molecules, have still not been identified. In addition, structural and

- 2 -

functional analyses of polypeptide products from many human genes have not been undertaken.

Accordingly, it is an object of the invention to identify novel polypeptides and nucleic acid molecules encoding the same which have diagnostic or  
5 therapeutic benefit.

#### Summary of the Invention

The present invention relates to novel FGF-like nucleic acid molecules and encoded polypeptides.

10 The invention provides for an isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence as set forth in SEQ ID NO: 1 or SEQ ID NO: 3;
- (b) the nucleotide sequence of the DNA insert in ATCC Deposit No.  
15 PTA-626;
- (c) a nucleotide sequence encoding the polypeptide as set forth in SEQ ID NO: 2 or SEQ ID NO: 4;
- (d) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a)-(c); and
- 20 (e) a nucleotide sequence complementary to any of (a)-(c).

The invention also provides for an isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence encoding a polypeptide that is at least about  
25 80 percent identical to the polypeptide as set forth in SEQ ID NO: 2 or SEQ ID NO: 4, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;

- 3 -

(b) a nucleotide sequence encoding an allelic variant or splice variant of the nucleotide sequence as set forth in SEQ ID NO: 1, SEQ ID NO: 3, or (a);

(c) a region of the nucleotide sequence of SEQ ID NO: 1, SEQ ID NO: 3, (a), or (b) encoding a polypeptide fragment of at least about 25 amino acid residues, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, has an oncogenic activity, or serves as an antigen for generating antibodies;

10 (d) a region of the nucleotide sequence of SEQ ID NO: 1, SEQ ID NO: 3, or any of (a) - (c) comprising a fragment of at least about 16 nucleotides;

(e) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (d); and

(f) a nucleotide sequence complementary to any of (a) - (d).

15

The invention further provides for an isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

(a) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one conservative amino acid substitution, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;

(b) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one amino acid insertion, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;



- 4 -

(c) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one amino acid deletion, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion  
5 from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;

(d) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 which has a carboxyl- and/or amino-terminal truncation, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth  
10 and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;

(e) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one modification selected from the group consisting of  
15 amino acid substitutions, amino acid insertions, amino acid deletions, carboxyl-terminal truncation, and amino-terminal truncation, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas  
20 chemotaxis, or has an oncogenic activity;

(f) a region of the nucleotide sequence of any of (a) - (e) comprising a fragment of at least about 16 nucleotides;

(g) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (f); and

25 (h) a nucleotide sequence complementary to any of (a) - (e).

The present invention provides for an isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

(a) the amino acid sequence as set forth in SEQ ID NO: 2 or SEQ ID  
30 NO: 4; and

- 5 -

(b) the amino acid sequence encoded by the DNA insert of ATCC Deposit No. PTA-626.

5 The invention also provides for an isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

(a) the amino acid sequence as set forth in either SEQ ID NO: 5 or SEQ ID NO: 6, optionally further comprising an amino-terminal methionine;

(b) an amino acid sequence for an ortholog of either SEQ ID NO: 2 or SEQ ID NO: 4, wherein the encoded polypeptide activates one or more FGF  
10 receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;

(c) an amino acid sequence that is at least about 80 percent identical to the amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 4, wherein the  
15 encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;

(d) a fragment of the amino acid sequence set forth in either SEQ ID  
20 NO: 2 or SEQ ID NO: 4 comprising at least about 25 amino acid residues, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, has an oncogenic activity, or serves as an antigen for  
25 generating antibodies; and

(e) an amino acid sequence for an allelic variant or splice variant of either the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 4; the amino acid sequence encoded by the DNA insert of ATCC Deposit No. PTA-626; (a), (b), or (c).

30

- 6 -

The invention further provides for an isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 4 with at least one conservative amino acid substitution, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;
- (b) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 4 with at least one amino acid insertion, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;
- (c) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 4 with at least one amino acid deletion, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;
- (d) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 4 which has a C- and/or N- terminal truncation, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity; and
- (e) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 4 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-terminal truncation, and N-terminal truncation, wherein the encoded polypeptide

- 7 -

activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity.

5

The invention also provides for an expression vector comprising the nucleic acid molecules as set forth above, host cells comprising the expression vectors of the invention, and a method of production of an FGF-like polypeptide comprising culturing the host cells and optionally isolating the polypeptide so  
10 produced.

A transgenic non-human animal comprising a nucleic acid molecule encoding an FGF-like polypeptide is also encompassed by the invention. The FGF-like nucleic acid molecules are introduced into the animal in a manner that allows expression and increased levels of an FGF-like polypeptide, which may  
15 include increased circulating levels. Alternatively, the FGF-like nucleic acid molecules are introduced into the animal in a manner that prevents expression of endogenous FGF-like polypeptide (*i.e.*, generates a transgenic animal possessing an FGF-like polypeptide gene knockout). The transgenic non-human animal is preferably a mammal, and more preferably a rodent, such as a rat or a mouse.  
20 Preferably the FGF-like transgene is expressed in the liver under the control of the apolipoprotein E promoter, or ubiquitously under the control of the beta actin promoter.

Also provided are derivatives of the FGF-like polypeptides of the invention, fusion polypeptides comprising the FGF-like polypeptides of the  
25 invention, and antibodies specifically binding the FGF-like polypeptides of the invention.

Compositions comprising the nucleotides or polypeptides of the invention and a carrier, adjuvant, solubilizer, stabilizer or anti-oxidant, or other pharmaceutically acceptable agent are also encompassed by the invention. The  
30 compositions may include pharmaceutical compositions comprising

- 8 -

therapeutically effective amounts of the nucleotides or polypeptides of the invention, and methods of using the polypeptides and nucleic acid molecules.

Surprisingly, FGF-like polypeptide appeared to be primarily expressed in the liver (Northern analysis) and pancreatic islets (*in situ* analysis), thereby distinguishing it from all other members of the FGF family. The present polypeptide, and its useful nucleic acid intermediates, may have utility, therefore, in differentiating liver cells or pancreatic islet cells from background. Further, given the localization of FGF-like polypeptide expression, the structural similarity of FGF-like polypeptide to members of the FGF family, and the likelihood that FGF-like polypeptide is secreted into the bloodstream where it may exert effects on distal sites, the present polypeptides may provide benefits in the stimulation of cells within or near the liver, regulation of intestinal cell activity, stimulation of cells within or near pancreatic islets, regulation of neuronal cells, stimulation or inhibition of angiogenesis, stimulation of epithelium or mesenchymal components of granulation tissue, stimulation of corneal epithelium, lens, or retinal tissue, regeneration of renal tubules, hematopoietic cell regulation, regulation of hair follicle growth, regulation of pulmonary epithelium, or stimulation of either epithelial, mesenchymal, hematopoietic, or neuronal cells or tissues, particularly as a therapeutic pharmaceutical composition.

FGF-like polypeptides may also be useful as growth or fat deposition inhibitors, and therefore may be useful in the treatment of excessive growth (for example, acromegaly), premature maturation, obesity or diabetes. Inhibitors – such as antibodies, binding proteins or small molecules – that interfere with the interaction of FGF-like polypeptides and their receptor(s) may be useful in stimulating body growth and maturation. Therefore, such inhibitors may be useful in treating short stature, delayed maturation, or other conditions generally associated with impairment of signaling of growth hormone or its mediator, insulin-like growth factor.

The FGF-like polypeptides and nucleic acid molecules of the invention, or agonists or antagonists of their biological activity, may be used for therapeutic or

- 9 -

diagnostic purposes to treat, prevent and/or detect a medical condition such as cirrhosis or other toxic insult of the liver; inflammatory bowel disease, mucositis, Crohn's disease, or other gastrointestinal abnormality; diabetes; obesity; neurodegenerative diseases; wounds; damage to the corneal epithelium, lens, or retinal tissue; damage to renal tubules as a result of acute tubular necrosis; hematopoietic cell reconstitution following chemotherapy; wasting syndromes (for example, cancer associated cachexia), multiple sclerosis, myopathies; short stature, delayed maturation, excessive growth (for example, acromegaly), premature maturation; alopecia; diseases or abnormalities of androgen target organs; infantile respiratory distress syndrome, bronchopulmonary dysplasia, acute respiratory distress syndrome, or other lung abnormalities; ~~or~~ tumors of the eye or other tissues; atherosclerosis; hypercholesterolemia; diabetes; obesity; stroke; osteoporosis; osteoarthritis; degenerative joint disease; muscle atrophy; sarcopenia; decreased lean body mass; baldness; wrinkles; increased fatigue; decreased stamina; decreased cardiac function; immune system dysfunction; cancer; Parkinson's disease; senile dementia; Alzheimer's disease; and decreased cognitive function. The invention provides for treating, preventing or ameliorating a disorder comprising administering to an animal an FGF-like polypeptide. The invention also provides for a method of diagnosing such a disorder or a susceptibility to such a disorder in an animal which includes both determining the presence or amount of expression of an FGF-like polypeptide and diagnosing such a disorder or a susceptibility to such a disorder based on the presence or amount of expression of an FGF-like polypeptide. The animal is preferably a mammal, and more preferably a human. The present invention also relates to methods for the manufacture of a medicament for the treatment of a disorder such as those mentioned above.

The invention also provides for the use of antibodies or other inhibitors of the binding of FGF-like polypeptide to its receptor for the treatment of the same diseases listed above, and for the treatment of tumors.



- 10 -

The invention also provides for a method of identifying a test molecule which binds to an FGF-like polypeptide wherein the method comprises contacting an FGF-like polypeptide with a test molecule and determining the extent of binding of the test molecule to the polypeptide. The method further comprises  
5 determining whether such test molecules are agonists or antagonists of an FGF-like polypeptide.

The invention also provides for a method of testing the impact of molecules on the expression of FGF-like polypeptide or on the activity of FGF-like polypeptide.

10 A method of regulating expression and modulating (*i.e.*, increasing or decreasing) levels of an FGF-like polypeptide are also encompassed by the invention. One method comprises administering to an animal a nucleic acid molecule encoding an FGF-like polypeptide. In another method, a nucleic acid molecule comprising elements that regulate expression of an FGF-like  
15 polypeptide may be administered. Examples of these methods include gene therapy and anti-sense therapy.

#### Description of the Figures

Figure 1 illustrates the nucleotide sequence of the murine FGF-like gene (SEQ ID  
20 NO: 3) and the deduced amino acid sequence of murine FGF-like protein (SEQ ID NO: 4);

Figures 2A-2B illustrate the nucleotide sequence of the human FGF-like gene (SEQ ID NO: 1) and the deduced amino acid sequence of human FGF-like protein  
25 (SEQ ID NO: 2);

Figures 3A-3D illustrate the amino acid sequence alignment of human FGF-like protein (hAgp-26257; SEQ ID NO: 2), murine FGF-like protein (mAgp-26257; SEQ ID NO: 4), human FGF-14 (Hfgf14; SEQ ID NO: 16), murine FGF-14  
30 (Mfgf14; SEQ ID NO: 26), human FGF-12 (Hfgf12; SEQ ID NO: 15), murine

- 11 -

FGF-13 (Mfgf13; SEQ ID NO: 25), human FGF-5 (Hfgf5; SEQ ID NO: 20),  
murine FGF-5 (Mfgf5; SEQ ID NO: 30), human FGF-6 (Hfgf6; SEQ ID NO: 21),  
murine FGF-6 (Mfgf6; SEQ ID NO: 31), human FGF-4 (Hfgf4; SEQ ID NO: 19),  
murine FGF-4 (Mfgf4; SEQ ID NO: 29), human FGF-3 (Hfgf3; SEQ ID NO: 18),  
5 murine FGF-3 (Mfgf3; SEQ ID NO: 28), human FGF-7 (Hfgf7; SEQ ID NO: 22),  
murine FGF-7 (Mfgf7; SEQ ID NO: 32), human FGF-9 (Hfgf9; SEQ ID NO: 23),  
murine FGF-9 (Mfgf9; SEQ ID NO: 33), human FGF-1 (Hfgf1; SEQ ID NO: 14),  
murine FGF-1 (Mfgf1; SEQ ID NO: 24), human FGF-2 (Hfgf2; SEQ ID NO: 17),  
murine FGF-2 (Mfgf2; SEQ ID NO: 27), and the resulting FGF consensus  
10 sequence (cons);

Figures 4A-4C illustrate the results of (A) a Northern blot analysis of murine  
FGF-like polypeptide expression, (B) a Northern blot analysis of human FGF-like  
polypeptide expression, and (C) a dot blot analysis of human FGF-like  
15 polypeptide expression.

#### Detailed Description of the Invention

The section headings herein are for organizational purposes only and are  
not to be construed as limiting the subject matter described therein. All references  
20 cited in this application are expressly incorporated by reference herein.

#### Definitions

The term "FGF-like nucleic acid molecule" refers to a nucleic acid  
molecule comprising or consisting essentially of a nucleotide sequence as set forth  
25 in SEQ ID NO: 1 or SEQ ID NO: 3, comprising or consisting essentially of a  
nucleotide sequence encoding the polypeptide as set forth in SEQ ID NO: 2 or  
SEQ ID NO: 4, comprising or consisting essentially of a nucleotide sequence of  
the DNA insert in ATCC Deposit No. PTA-626, or nucleic acid molecules related  
thereto.

- 12 -

Related nucleic acid molecules comprise or consist essentially of a nucleotide sequence that is about 80 percent identical to the nucleotide sequence as shown in SEQ ID NO: 1 or SEQ ID NO: 3, or comprise or consist essentially of a nucleotide sequence encoding a polypeptide that is about 80 percent identical to the polypeptide as set forth in SEQ ID NO: 2 or SEQ ID NO: 4. In preferred embodiments, the nucleotide sequences are about 85 percent, or about 90 percent, or about 95 percent, or about 96, 97, 98, or 99 percent identical to the nucleotide sequence as shown in SEQ ID NO: 1 or SEQ ID NO: 3, or the nucleotide sequences encode a polypeptide that is about 85 percent, or about 90 percent, or about 95 percent, or about 96, 97, 98, or 99 percent identical to the polypeptide sequence as set forth in SEQ ID NO: 2 or SEQ ID NO: 4. Related nucleic acid molecules also include fragments of the above FGF-like nucleic acid molecules which are at least about 16 contiguous nucleotides, or about 18, or about 20, or about 25, or about 50, or about 75, or about 100, or greater than about 100 contiguous nucleotides. Related nucleic acid molecules also include fragments of the above FGF-like nucleic acid molecules which encode a polypeptide of at least about 25 amino acid residues, or about 50, or about 75, or about 100, or greater than about 100 amino acid residues.

Related nucleic acid molecules also include a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 4 with at least one conservative amino acid substitution and wherein the polypeptide retains at least one FGF-like polypeptide activity, or a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 4 with at least one amino acid insertion and wherein the polypeptide retains at least one FGF-like polypeptide activity, or a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 4 with at least one amino acid deletion and wherein the polypeptide retains at least one FGF-like polypeptide activity. Related nucleic acid molecules further include a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 4 which has a C- and/or N- terminal truncation and wherein the polypeptide retains

- 13 -

at least one FGF-like polypeptide activity. Related nucleic acid molecules also include a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 4 with combinations of modifications selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-terminal truncations, and N-terminal truncations and wherein the polypeptide retains at least one FGF-like polypeptide activity.

Related FGF-like nucleic acid molecules include those molecules that comprise nucleotide sequences which hybridize under moderately or highly stringent conditions as defined herein with the complements of any of the above nucleic acid molecules. In preferred embodiments, the related nucleic acid molecules comprise sequences which hybridize under moderately or highly stringent conditions with the sequence as shown in SEQ ID NO: 1 or SEQ ID NO: 3, or with a molecule encoding a polypeptide, which polypeptide comprises the sequence as shown in SEQ ID NO: 2 or SEQ ID NO: 4, or with a nucleic acid fragment as defined above, or with a nucleic acid fragment encoding a polypeptide as defined above. It is also understood that related nucleic acid molecules include allelic or splice variants of any of the above nucleic acids, and include sequences which are complementary to any of the above nucleotide sequences.

The term "isolated nucleic acid molecule" refers to a nucleic acid molecule of the invention that is free from at least one contaminating nucleic acid molecule with which it is naturally associated, and preferably substantially free from any other contaminating mammalian nucleic acid molecules which would interfere with its use in protein production or its therapeutic or diagnostic use.

The term "allelic variant" refers to one of several possible naturally occurring alternate forms of a gene occupying a given locus on a chromosome of an organism or a population of organisms.

The term "splice variant" refers to a nucleic acid molecule, usually RNA, which is generated by alternative processing of intron sequences in an RNA transcript.

- 14 -

The term "expression vector" refers to a vector that is suitable for propagation in a host cell and contains nucleic acid sequences that direct and/or control the expression of inserted heterologous nucleic acid sequences. Expression includes, but is not limited to, processes such as transcription, translation, and RNA splicing, if introns are present.

The term "highly stringency conditions" refers to those conditions that (1) employ low ionic strength reagents and high temperature for washing, for example, 0.015 M NaCl/0.0015 M sodium citrate/0.1% NaDodSO<sub>4</sub> (SDS) at 50°C, or (2) employ during hybridization a denaturing agent such as formamide, for example, 50% (vol/vol) formamide with 0.1% bovine serum albumin, 0.2% Ficoll, 0.1% polyvinylpyrrolidone, 50 mM sodium phosphate buffer (pH 6.5), 750 mM NaCl, and 75 mM sodium citrate at 42°C. Another example is the use of 50% formamide, 5X SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5X Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2X SSC and 0.1% SDS.

The term "moderately stringency conditions" refers to conditions which generally include the use of a washing solution and hybridization conditions (e.g., temperature, ionic strength, and percentage of SDS) less stringent than described above. An example of moderately stringent conditions are conditions such as overnight incubation at 37°C in a solution comprising 20% formamide, 5X SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5X Denhardt's solution, 10% dextran sulfate, and 20 µl/ml denatured sheared salmon sperm DNA, followed by washing in 1X SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

In certain preferred embodiments, where oligonucleotide probes are used to screen cDNA or genomic libraries, high stringency conditions are used which depend upon the melting temperature ( $T_m$ ) of oligonucleotide probes to target

- 15 -

sequences. The  $T_m$  may be estimated using the following formula (Bolton et al., *Proc. Natl. Acad. Sci. U.S.A.* 48:1390 (1962)):

$$T_m = 81.5 - 16.6 (\log[\text{Na}^+]) + 0.41(\%G+C) - (600/N)$$

5

wherein  $[\text{Na}^+]$  is the sodium ion concentration in the hybridization (or washing) solution;

$\%G+C$  is guanine and cytosine content in the oligonucleotide probe; and

$N$  is the probe length in nucleotides.

10

An example of a high stringency solution is 6X SSC and 0.05% sodium pyrophosphate at a temperature of 35-63°C, depending on the length of the oligonucleotide probe. For example, according to certain embodiments, 14 base pair probes are washed at 35-40°C, 17 base pair probes are washed at 45-50°C, 20  
15 base pair probes are washed at 52-57°C, and 23 base pair probes are washed at 57-63°C. The temperature can be increased 2-3°C where the background non-specific binding appears high. A second high stringency solution utilizes tetramethylammonium chloride (TMAC) for washing oligonucleotide probes. One stringent washing solution is 3 M TMAC, 50 mM Tris-HCl, pH 8.0, and  
20 0.2% SDS. The washing temperature using this solution is a function of the length of the probe. For example, 14 base pair probes are washed at 35-40°C, 17 base pair probes are washed at about 45-50°C, 20 base pair probes are washed at 52-57°C, and 23 base pair probes are washed at 57-63°C.

The term "FGF-like polypeptides" refers to a polypeptide comprising the  
25 amino acid sequence of SEQ ID NO: 2 or SEQ ID NO: 4, and related polypeptides described herein. Related polypeptides include: allelic variants; splice variants; fragments; derivatives; substitution, deletion, and insertion variants; fusion polypeptides; and orthologs. FGF-like polypeptides may be mature polypeptides, as defined herein, and may or may not have an amino terminal methionine  
30 residue, depending on the method by which they are prepared.



- 16 -

The term "FGF-like polypeptide fragment" refers to a peptide or polypeptide that comprises less than the full length amino acid sequence of an FGF-like polypeptide as set forth in SEQ ID NO: 2 or SEQ ID NO: 4. Such a fragment may arise, for example, from a truncation at the amino terminus, a  
5 truncation at the carboxyl terminus, and/or an internal deletion of a residue(s) from the amino acid sequence. FGF-like fragments may result from alternative RNA splicing or from *in vivo* protease activity.

The term "FGF-like polypeptide variants" refers to FGF-like polypeptides comprising amino acid sequences which contain one or more amino acid sequence  
10 substitutions, deletions, and/or additions as compared to the FGF-like polypeptide amino acid sequence set forth in SEQ ID NO: 2 or SEQ ID NO: 4. Variants may be naturally occurring or artificially constructed. Such FGF-like polypeptide variants may be prepared from the corresponding nucleic acid molecules encoding said variants, which have a DNA sequence that varies accordingly from the DNA  
15 sequences for wild type FGF-like polypeptides as set forth in SEQ ID NO: 1 or SEQ ID NO: 3.

The term "FGF-like fusion polypeptide" refers to a fusion of an FGF-like polypeptide, fragment, variant, or derivative thereof, with a heterologous peptide or polypeptide.

20 The term "FGF-like polypeptide derivatives" refers to FGF-like polypeptides, variants, or fragments thereof, that have been chemically modified, as for example, by covalent attachment of one or more polymers, including, but not limited to, water soluble polymers, N-linked or O-linked carbohydrates, sugars, phosphates, and/or other such molecules. The derivatives are modified in  
25 a manner that is different from naturally occurring FGF-like polypeptide, either in the type or location of the molecules attached to the polypeptide. Derivatives further include the deletion of one or more chemical groups naturally attached to the FGF-like polypeptide.

The terms "biologically active FGF-like polypeptides," "biologically  
30 active FGF-like polypeptide fragments," "biologically active FGF-like

- 17 -

polypeptide variants,” and “biologically active FGF-like polypeptide derivatives” refer to FGF-like polypeptides having at least one activity characteristic of an FGF-like polypeptide, such as stimulation of cells within or near the liver, regulation of intestinal cell activity, stimulation of cells within or near pancreatic islets, regulation of neuronal cells, stimulation or inhibition of angiogenesis, stimulation of epithelium or mesenchymal components of granulation tissue, stimulation of corneal epithelium, lens, or retinal tissue, regeneration of renal tubules, hematopoietic cell regulation, regulation of hair follicle growth, regulation of pulmonary epithelium, or stimulation of either epithelial, mesenchymal, hematopoietic, or neuronal cells or tissues. In general, FGF-like polypeptides, and variants, fragments and derivatives thereof, will have at least one activity characteristic of an FGF-like polypeptide such as those activities listed above. In addition, an FGF-like polypeptide may be active as an immunogen (*i.e.*, the polypeptide contains at least one epitope to which antibodies may be raised).

“Naturally occurring” when used in connection with biological materials such as nucleic acid molecules, polypeptides, host cells, and the like, refers to that which are found in nature and not manipulated by a human being.

The term “isolated polypeptide” refers to a polypeptide of the invention that is free from at least one contaminating polypeptide that is found in its natural environment, and preferably substantially free from any other contaminating mammalian polypeptides which would interfere with its therapeutic or diagnostic use.

The term “ortholog” refers to a polypeptide that corresponds to a polypeptide identified from a different species. For example, murine and human FGF-like polypeptides are considered orthologs of one another.

The term “mature FGF-like polypeptide” refers to a polypeptide lacking a leader sequence and may also include other modifications of a polypeptide such as proteolytic processing of the amino terminus (with or without a leader sequence) and/or the carboxyl terminus, cleavage of a smaller polypeptide from a larger

- 18 -

precursor, N-linked and/or O-linked glycosylation, and other post-translational modifications understood by those with skill in the art.

The terms "effective amount" and "therapeutically effective amount" refer to the amount of a FGF-like polypeptide that is useful or necessary to support an observable level of one or more biological activities of the FGF-like polypeptides as set forth above.

#### Relatedness of Nucleic Acid Molecules and/or Polypeptides

The term "identity," as known in the art, refers to a relationship between the sequences of two or more polypeptide molecules or two or more nucleic acid molecules, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or nucleic acid molecule sequences, as the case may be, as determined by the match between strings of nucleotide or amino acid sequences. "Identity" measures the percent of identical matches between two or more sequences with gap alignments addressed by a particular mathematical model of computer programs (*i.e.*, "algorithms").

The term "similarity" is a related concept, but in contrast to "identity," refers to a measure of similarity which includes both identical matches and conservative substitution matches. Since conservative substitutions apply to polypeptides and not nucleic acid molecules, similarity only deals with polypeptide sequence comparisons. If two polypeptide sequences have, for example, 10 out of 20 identical amino acids, and the remainder are all non-conservative substitutions, then the percent identity and similarity would both be 50%. If in the same example, there are 5 more positions where there are conservative substitutions, then the percent identity remains 50%, but the percent similarity would be 75% (15 out of 20). Therefore, in cases where there are conservative substitutions, the degree of similarity between two polypeptide sequences will be higher than the percent identity between those two sequences.

The term "conservative amino acid substitution" refers to a substitution of a native amino acid residue with a nonnative residue such that there is little or no

- 19 -

effect on the polarity or charge of the amino acid residue at that position. For example, a conservative substitution results from the replacement of a non-polar residue in a polypeptide with any other non-polar residue. Furthermore, any native residue in the polypeptide may also be substituted with alanine, as has been

5 previously described for "alanine scanning mutagenesis" (Cunnigham et al., *Science* 244:1081-85 (1989)). General rules for conservative amino acid substitutions are set forth in Table I.

Table I

## 10 Conservative Amino Acid Substitutions

Original Residues	Exemplary Substitutions	Preferred Substitutions
Ala	Val, Leu, Ile	Val
Arg	Lys, Gln, Asn	Lys
Asn	Gln, His, Lys, Arg	Gln
Asp	Glu	Glu
Cys	Ser	Ser
Gln	Asn	Asn
Glu	Asp	Asp
Gly	Pro, Ala	Ala
His	Asn, Gln, Lys, Arg	Arg
Ile	Leu, Val, Met, Ala, Phe, Norleucine	Leu
Leu	Norleucine, Ile, Val, Met, Ala, Phe	Ile
Lys	Arg, Gln, Asn	Arg
Met	Leu, Phe, Ile	Leu
Phe	Leu, Val, Ile, Ala, Tyr	Leu
Pro	Ala	Ala

- 20 -

Ser	Thr	Thr
Thr	Ser	Ser
Trp	Tyr, Phe	Tyr
Tyr	Trp, Phe, Thr, Ser	Phe
Val	Ile, Met, Leu, Phe,	Leu
	Ala, Norleucine	

Conservative amino acid substitutions also encompass non-naturally occurring amino acid residues that are typically incorporated by chemical peptide synthesis rather than by synthesis in biological systems. These include peptidomimetics,  
5 and other reversed or inverted forms of amino acid moieties.

Conservative modifications to the amino acid sequence (and the corresponding modifications to the encoding nucleotides) are expected to produce FGF-like polypeptide having functional and chemical characteristics similar to those of naturally occurring FGF-like polypeptide. In contrast, substantial  
10 modifications in the functional and/or chemical characteristics of FGF-like polypeptide may be accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the molecular backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site,  
15 or (c) the bulk of the side chain. Naturally occurring residues may be divided into groups based on common side chain properties:

- 1) hydrophobic: norleucine, Met, Ala, Val, Leu, Ile;
- 2) neutral hydrophilic: Cys, Ser, Thr;
- 3) acidic: Asp, Glu;
- 20 4) basic: Asn, Gln, His, Lys, Arg;
- 5) residues that influence chain orientation: Gly, Pro; and
- 6) aromatic: Trp, Tyr, Phe.

Non-conservative substitutions may involve the exchange of a member of one of these classes for a member from another class. Such substituted residues

- 21 -

may be introduced into regions of the human FGF-like molecule that are homologous with non-human FGF-like polypeptide, or into the non-homologous regions of the molecule.

Identity and similarity of related nucleic acid molecules and polypeptides  
5 can be readily calculated by known methods, including but not limited to those described in *Computational Molecular Biology* (A.M. Lesk, ed., Oxford University Press 1988); *Biocomputing: Informatics and Genome Projects* (D.W. Smith, ed., Academic Press 1993); *Computer Analysis of Sequence Data* (Part 1, A.M. Griffin and H.G. Griffin, eds., Humana Press 1994); G. von Heinle,  
10 *Sequence Analysis in Molecular Biology* (Academic Press 1987); *Sequence Analysis Primer* (M. Gribskov and J. Devereux, eds., M. Stockton Press 1991); and Carillo et al., *SIAM J. Applied Math.* 48:1073 (1988).

Preferred methods to determine identity and/or similarity are designed to give the largest match between the sequences tested. Methods to determine  
15 identity and similarity are codified in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, the GCG program package, including GAP (Devereux et al., *Nuc. Acids Res.* 12:387 (1984); Genetics Computer Group, University of Wisconsin, Madison, WI), BLASTP, BLASTN,  
20 and FASTA (Atschul et al., *J. Mol. Biol.* 215:403-10 (1990)). The BLAST X program is publicly available from the National Center for Biotechnology Information (NCBI) and other sources (Altschul et al., *BLAST Manual* (NCB NLM NIH, Bethesda, MD); Altschul et al., 1990, *supra*). The well-known Smith Waterman algorithm may also be used to determine identity.

25 By way of example, using the computer algorithm GAP (Genetics Computer Group), two polypeptides for which the percent sequence identity is to be determined are aligned for optimal matching of their respective amino acids (the "matched span," as determined by the algorithm). A gap opening penalty (which is calculated as 3X the average diagonal; the "average diagonal" is the  
30 average of the diagonal of the comparison matrix being used; the "diagonal" is the



- 22 -

score or number assigned to each perfect amino acid match by the particular comparison matrix) and a gap extension penalty (which is usually 0.1X the gap opening penalty), as well as a comparison matrix such as PAM 250 or BLOSUM 62 are used in conjunction with the algorithm. A standard comparison matrix (see  
5 Dayhoff et al., 5 *Atlas of Protein Sequence and Structure* (Supp. 3 1978) for the PAM250 comparison matrix; see Henikoff et al., *Proc. Natl. Acad. Sci USA* 89:10915-19 (1992) for the BLOSUM 62 comparison matrix) is also used by the algorithm.

Preferred parameters for polypeptide sequence comparison include the  
10 following:

Algorithm: Needleman and Wunsch, *J. Mol. Biol.* 48:443-53 (1970)  
Comparison matrix: BLOSUM 62 from Henikoff et al., *Proc. Natl. Acad. Sci. U.S.A.* 89:10915-19 (1992)  
15 Gap Penalty: 12  
Gap Length Penalty: 4  
Threshold of Similarity: 0

The GAP program is useful with the above parameters. The aforementioned  
20 parameters are the default parameters for polypeptide comparisons (along with no penalty for end gaps) using the GAP algorithm.

Preferred parameters for nucleic acid molecule sequence comparison include the following:

25 Algorithm: Needleman et al., *J. Mol Biol.* 48:443-53 (1970)  
Comparison matrix: matches = +10, mismatch = 0  
Gap Penalty: 50  
Gap Length Penalty: 3

- 23 -

The GAP program is also useful with the above parameters. The aforementioned parameters are the default parameters for nucleic acid molecule comparisons.

Other exemplary algorithms, gap opening penalties, gap extension penalties, comparison matrices, thresholds of similarity, etc. may be used by those of skill in the art, including those set forth in the Program Manual, Wisconsin Package, Version 9, September, 1997. The particular choices to be made will depend on the specific comparison to be made, such as DNA to DNA, protein to protein, protein to DNA; and additionally, whether the comparison is between given pairs of sequences (in which case GAP or BestFit are generally preferred) or between one sequence and a large database of sequences (in which case FASTA or BLASTA are preferred).

Sequence analysis of an isolated mouse cDNA (murine FGF-like protein; SEQ ID NO: 3) indicated that it encoded a novel member of the FGF family of proteins. The murine FGF-like gene comprises a 630 bp open reading frame encoding a protein of 210 amino acids (Figure 1). The murine sequence was used to identify the human FGF-like ortholog. Sequence analysis of four human FGF-like polypeptide cDNA clones indicated that the human FGF-like gene comprises a 627 bp open reading frame encoding a protein of 209 amino acids (Figures 2A-2B).

Figures 3A-3D illustrate the amino acid sequence alignment of human FGF-like protein, murine FGF-like protein, and other members of the FGF family. Computer analysis of the predicted murine FGF-like polypeptide, using the FASTA program of the Swissprot database, indicated that the protein was most closely related to murine FGF-6, FGF-15, and FGF-4. Using the GAP program, murine FGF-like polypeptide was found to be 32% identical to murine FGF-6 and 28% identical to murine FGF-4. Computer analysis also indicated that the murine FGF-like polypeptide, similar to FGF-6, FGF-4, and FGF-15 but in contrast to FGF-1 and FGF-2, possessed a potential signal peptide at its amino terminus. The murine FGF-like polypeptide is 79% identical to the human FGF-like protein.

### Nucleic Acid Molecules

Recombinant DNA methods used herein are generally those set forth in Sambrook et al., *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1989) and/or *Current Protocols in Molecular Biology* (Ausubel  
5 et al., eds., Green Publishers Inc. and Wiley and Sons 1994).

The invention provides for nucleic acid molecules as described herein and methods for obtaining the molecules. A gene or cDNA encoding an FGF-like polypeptide or fragment thereof may be obtained by hybridization screening of a genomic or cDNA library, or by PCR amplification. Probes or primers useful for  
10 screening a library by hybridization can be generated based on sequence information for other known genes or gene fragments from the same or a related family of genes, such as, for example, conserved motifs. In addition, where a gene encoding FGF-like polypeptide has been identified from one species, all or a portion of that gene may be used as a probe to identify corresponding genes from  
15 other species (orthologs) or related genes from the same species (homologs). The probes or primers may be used to screen cDNA libraries from various tissue sources believed to express the FGF-like gene. In addition, part or all of a nucleic acid molecule having the sequence as set forth in SEQ ID NO: 1 or SEQ ID NO: 3 may be used to screen a genomic library to identify and isolate a gene encoding  
20 FGF-like polypeptide. Typically, conditions of moderate or high stringency will be employed for screening to minimize the number of false positives obtained from the screen.

Nucleic acid molecules encoding FGF-like polypeptides may also be identified by expression cloning which employs detection of positive clones based  
25 upon a property of the expressed protein. Typically, nucleic acid libraries are screened by binding an antibody or other binding partner (e.g., receptor or ligand) to cloned proteins which are expressed and displayed on the host cell surface. The antibody or binding partner is modified with a detectable label to identify those cells expressing the desired clone.

- 25 -

Another means of preparing a nucleic acid molecule encoding an FGF-like polypeptide or fragment thereof is chemical synthesis using methods well known to the skilled artisan such as those described by Engels et al., *Angew. Chem. Intl. Ed.* 28:716-34 (1989). These methods include, *inter alia*, the phosphotriester, 5 phosphoramidite, and H-phosphonate methods for nucleic acid synthesis. A preferred method for such chemical synthesis is polymer-supported synthesis using standard phosphoramidite chemistry. Typically, the DNA encoding the FGF-like polypeptide will be several hundred nucleotides in length. Nucleic acids larger than about 100 nucleotides can be synthesized as several fragments using 10 these methods. The fragments can then be ligated together to form the full-length FGF-like polypeptide. Usually, the DNA fragment encoding the amino terminus of the polypeptide will have an ATG, which encodes a methionine residue. This methionine may or may not be present on the mature form of the FGF-like polypeptide, depending on whether the polypeptide produced in the host cell is 15 designed to be secreted from that cell.

In some cases, it may be desirable to prepare nucleic acid molecules encoding FGF-like polypeptide variants. Nucleic acid molecules encoding variants may be produced using site directed mutagenesis, PCR amplification, or other appropriate methods, where the primer(s) have the desired point mutations 20 (*see* Sambrook et al., *supra*, and Ausubel et al., *supra*, for descriptions of mutagenesis techniques). Chemical synthesis using methods described by Engels et al., *supra*, may also be used to prepare such variants. Other methods known to the skilled artisan may be used as well.

In certain embodiments, nucleic acid variants contain codons which have 25 been altered for optimal expression of an FGF-like polypeptide in a given host cell. Particular codon alterations will depend upon the FGF-like polypeptide and host cell selected for expression. Such "codon optimization" can be carried out by a variety of methods, for example, by selecting codons which are preferred for use in highly expressed genes in a given host cell. Computer algorithms which 30 incorporate codon frequency tables such as "Ecohigh.\_Cod" for codon preference

- 26 -

of highly expressed bacterial genes may be used and are provided by the University of Wisconsin Package Version 9.0, Genetics Computer Group, Madison, WI. Other useful codon frequency tables include "Celegans\_high.cod," "Celegans\_low.cod," "Drosophila\_high.cod," "Human\_high.cod,"  
5 "Maize\_high.cod," and "Yeast\_high.cod."

In other embodiments, nucleic acid molecules encode FGF-like variants with conservative amino acid substitutions as defined above, FGF-like variants comprising an addition and/or a deletion of one or more N-linked or O-linked glycosylation sites, FGF-like variants having deletions and/or substitutions of one  
10 or more cysteine residues, or FGF-like polypeptide fragments as described above. In addition, nucleic acid molecules may encode any combination of FGF-like variants, fragments, and fusion polypeptides described herein.

#### Vectors and Host Cells

15 A nucleic acid molecule encoding an FGF-like polypeptide is inserted into an appropriate expression vector using standard ligation techniques. The vector is typically selected to be functional in the particular host cell employed (*i.e.*, the vector is compatible with the host cell machinery such that amplification of the gene and/or expression of the gene can occur). A nucleic acid molecule encoding  
20 an FGF-like polypeptide may be amplified/expressed in prokaryotic, yeast, insect (baculovirus systems) and/or eukaryotic host cells. Selection of the host cell will depend in part on whether an FGF-like polypeptide is to be post-translationally modified (*e.g.*, glycosylated and/or phosphorylated). If so, yeast, insect, or mammalian host cells are preferable. For a review of expression vectors, see  
25 *Meth. Enz.* (D.V. Goeddel, ed., Academic Press 1990).

Typically, expression vectors used in any of the host cells will contain sequences for plasmid maintenance and for cloning and expression of exogenous nucleotide sequences. Such sequences, collectively referred to as "flanking sequences" in certain embodiments will typically include one or more of the  
30 following nucleotides: a promoter, one or more enhancer sequences, an origin of

- 27 -

replication, a transcriptional termination sequence, a complete intron sequence containing a donor and acceptor splice site, a leader sequence for secretion, a ribosome binding site, a polyadenylation sequence, a polylinker region for inserting the nucleic acid encoding the polypeptide to be expressed, and a  
5 selectable marker element. Each of these sequences is discussed below.

Optionally, the vector may contain a "tag" sequence, *i.e.*, an oligonucleotide molecule located at the 5' or 3' end of the FGF-like polypeptide coding sequence; the oligonucleotide molecule encodes polyHis (such as hexaHis), or other "tag" such as FLAG, HA (hemagglutinin Influenza virus) or *myc*  
10 for which commercially available antibodies exist. This tag is typically fused to the polypeptide upon expression of the polypeptide, and can serve as a means for affinity purification of the FGF-like polypeptide from the host cell. Affinity purification can be accomplished, for example, by column chromatography using antibodies against the tag as an affinity matrix. Optionally, the tag can  
15 subsequently be removed from the purified FGF-like polypeptide by various means such as using certain peptidases for cleavage.

Flanking sequences may be homologous (*i.e.*, from the same species and/or strain as the host cell), heterologous (*i.e.*, from a species other than the host cell species or strain), hybrid (*i.e.*, a combination of flanking sequences from more  
20 than one source), or synthetic, or native sequences which normally function to regulate FGF-like expression. As such, the source of flanking sequences may be any prokaryotic or eukaryotic organism, any vertebrate or invertebrate organism, or any plant, provided that the flanking sequences is functional in, and can be activated by, the host cell machinery.

25 The flanking sequences useful in the vectors of this invention may be obtained by any of several methods well known in the art. Typically, flanking sequences useful herein other than the FGF-like gene flanking sequences will have been previously identified by mapping and/or by restriction endonuclease digestion and can thus be isolated from the proper tissue source using the  
30 appropriate restriction endonucleases. In some cases, the full nucleotide sequence



- 28 -

of one or more flanking sequence may be known. Here, the flanking sequence may be synthesized using the methods described above for nucleic acid synthesis or cloning.

Where all or only a portion of the flanking sequence is known, it may be  
5 obtained using PCR and/or by screening a genomic library with suitable oligonucleotide and/or flanking sequence fragments from the same or another species.

Where the flanking sequence is not known, a fragment of DNA containing a flanking sequence may be isolated from a larger piece of DNA that may contain,  
10 for example, a coding sequence or even another gene or genes. Isolation may be accomplished by restriction endonuclease digestion to produce the proper DNA fragment followed by isolation using agarose gel purification, Qiagen® (Valencia, CA) column chromatography, or other methods known to the skilled artisan. Selection of suitable enzymes to accomplish this purpose will be readily apparent  
15 to one of ordinary skill in the art.

An origin of replication is typically a part of prokaryotic expression vectors purchased commercially, and aids in the amplification of the vector in a host cell. Amplification of the vector to a certain copy number can, in some cases, be important for optimal expression of the FGF-like polypeptide. If the  
20 vector of choice does not contain an origin of replication site, one may be chemically synthesized based on a known sequence, and ligated into the vector.

The origin of replication from the plasmid pBR322 (Product No. 303-3s, New England Biolabs, Beverly, MA) is suitable for most Gram-negative bacteria and various origins (*e.g.*, SV40, polyoma, adenovirus, vesicular stomatitis virus  
25 (VSV) or papillomaviruses such as HPV or BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors (for example, the SV40 origin is often used only because it contains the early promoter).

A transcription termination sequence is typically located 3' of the end of a  
30 polypeptide coding regions and serves to terminate transcription. Usually, a

- 29 -

transcription termination sequence in prokaryotic cells is a G-C rich fragment followed by a poly-T sequence. While the sequence is easily cloned from a library or even purchased commercially as part of a vector, it can also be readily synthesized using methods for nucleic acid synthesis such as those described  
5 above.

A selectable marker gene element encodes a protein necessary for the survival and growth of a host cell grown in a selective culture medium. Typical selection marker genes encode proteins that (a) confer resistance to antibiotics or other toxins, for example, ampicillin, tetracycline, or kanamycin for prokaryotic  
10 host cells, (b) complement auxotrophic deficiencies of the cell; or (c) supply critical nutrients not available from complex media. Preferred selectable markers are the kanamycin resistance gene, the ampicillin resistance gene, and the tetracycline resistance gene. A neomycin resistance gene may also be used for selection in prokaryotic and eukaryotic host cells.

15 Other selection genes may be used to amplify the gene that will be expressed. Amplification is the process wherein genes that are in greater demand for the production of a protein critical for growth are reiterated in tandem within the chromosomes of successive generations of recombinant cells. Examples of suitable selectable markers for mammalian cells include dihydrofolate reductase  
20 (DHFR) and thymidine kinase. The mammalian cell transformants are placed under selection pressure that only the transformants are uniquely adapted to survive by virtue of the marker present in the vector. Selection pressure is imposed by culturing the transformed cells under conditions in which the concentration of selection agent in the medium is successively changed, thereby  
25 leading to amplification of both the selection gene and the DNA that encodes FGF-like polypeptide. As a result, increased quantities of FGF-like polypeptide are synthesized from the amplified DNA.

A ribosome binding site is usually necessary for translation initiation of mRNA and is characterized by a Shine-Dalgarno sequence (prokaryotes) or a  
30 Kozak sequence (eukaryotes). The element is typically located 3' to the promoter

- 30 -

and 5' to the coding sequence of the FGF-like polypeptide to be expressed. The Shine-Dalgarno sequence is varied but is typically a polypurine (*i.e.*, having a high A-G content). Many Shine-Dalgarno sequences have been identified, each of which can be readily synthesized using methods set forth above and used in a  
5 prokaryotic vector.

A leader, or signal, sequence may be used to direct an FGF-like polypeptide out of the host cell. Typically, the signal sequence is positioned in the coding region of the FGF-like nucleic acid molecule, or directly at the 5' end of the FGF-like polypeptide coding region. Many signal sequences have been  
10 identified, and any of them that are functional in the selected host cell may be used in conjunction with the FGF-like gene or cDNA. Therefore, a signal sequence may be homologous (naturally occurring) or heterologous to the FGF-like gene or cDNA. Additionally, a signal sequence may be chemically synthesized using methods set forth above. In most cases, secretion of an FGF-  
15 like polypeptide from the host cell via the presence of a signal peptide will result in the removal of the signal peptide from the FGF-like polypeptide. The signal sequence may be a component of the vector, or it may be a part of FGF-like DNA that is inserted into the vector.

Included within the scope of this invention is the native FGF-like signal  
20 sequence joined to an FGF-like coding region and a heterologous signal sequence joined to an FGF-like coding region. The heterologous signal sequence selected should be one that is recognized and processed, *i.e.*, cleaved by a signal peptidase, by the host cell. For prokaryotic host cells that do not recognize and process the native FGF-like signal sequence, the signal sequence is substituted by a  
25 prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, or heat-stable enterotoxin II leaders. For yeast secretion, the native FGF-like signal sequence may be substituted by the yeast invertase, alpha factor, or acid phosphatase leaders. In mammalian cell expression the native signal sequence is satisfactory, although other mammalian signal  
30 sequences may be suitable.

- 31 -

In some cases, such as where glycosylation is desired in a eukaryotic host cell expression system, one may manipulate the various presequences to improve glycosylation or yield. For example, one may alter the peptidase cleavage site of a particular signal peptide, or add presequences, which also may affect glycosylation. The final protein product may have, in the -1 position (relative to the first amino acid of the mature protein) one or more additional amino acids incident to expression, which may not have been totally removed. For example, the final protein product may have one or two amino acid found in the peptidase cleavage site, attached to the N-terminus. Alternatively, use of some enzyme cleavage sites may result in a slightly truncated form of the desired FGF-like polypeptide, if the enzyme cuts at such area within the mature polypeptide.

In many cases, transcription of a nucleic acid molecule is increased by the presence of one or more introns in the vector; this is particularly true where a polypeptide is produced in eukaryotic host cells, especially mammalian host cells. The introns used may be naturally occurring within the FGF-like gene especially where the gene used is a full-length genomic sequence or a fragment thereof. Where the intron is not naturally occurring within the gene (as for most cDNAs), the intron may be obtained from another source. The position of the intron with respect to flanking sequences and the FGF-like gene is generally important, as the intron must be transcribed to be effective. Thus, when an FGF-like cDNA molecule is being expressed, the preferred position for the intron is 3' to the transcription start site and 5' to the poly-A transcription termination sequence. Preferably, the intron or introns will be located on one side or the other (*i.e.*, 5' or 3') of the cDNA such that it does not interrupt the coding sequence. Any intron from any source, including any viral, prokaryotic and eukaryotic (plant or animal) organisms, may be used to practice this invention, provided that it is compatible with the host cell into which it is inserted. Also included herein are synthetic introns. Optionally, more than one intron may be used in the vector.

The expression and cloning vectors of the present invention will typically contain a promoter that is recognized by the host organism and operably linked to

- 32 -

the molecule encoding the FGF-like protein. Promoters are untranslated sequences located upstream (*i.e.*, 5') to the start codon of a structural gene (generally within about 100 to 1000 bp) that control the transcription and translation of the structural gene. Promoters are conventionally grouped into one of two classes: inducible promoters and constitutive promoters. Inducible promoters initiate increased levels of transcription from DNA under their control in response to some change in culture conditions, such as the presence or absence of a nutrient or a change in temperature. A large number of promoters, recognized by a variety of potential host cells, are well known. These promoters are operably linked to the DNA encoding FGF-like polypeptide by removing the promoter from the source DNA by restriction enzyme digestion and inserting the desired promoter sequence into the vector. The native FGF-like promoter sequence may be used to direct amplification and/or expression of FGF-like DNA. A heterologous promoter is preferred, however, if it permits greater transcription and higher yields of the expressed protein as compared to the native promoter, and if it is compatible with the host cell system that has been selected for use.

Promoters suitable for use with prokaryotic hosts include the beta-lactamase and lactose promoter systems; alkaline phosphatase, a tryptophan (*trp*) promoter system; and hybrid promoters such as the *tac* promoter. Other known bacterial promoters are also suitable. Their sequences have been published, thereby enabling one skilled in the art to ligate them to the desired DNA sequence, using linkers or adapters as needed to supply any required restriction sites.

Suitable promoters for use with yeast hosts are also well known in the art. Yeast enhancers are advantageously used with yeast promoters. Suitable promoters for use with mammalian host cells are well known and include those obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and most preferably Simian Virus 40 (SV40). Other suitable mammalian promoters include heterologous mammalian promoters, for example, heat-shock promoters and the actin promoter.



- 33 -

Additional promoters which may be of interest in controlling FGF-like gene expression include, but are not limited to: the SV40 early promoter region (Bernoist and Chambon, *Nature* 290:304-10 (1981)); the CMV promoter; the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto, et al., *Cell* 22:787-97 (1980)); the herpes thymidine kinase promoter (Wagner et al., *Proc. Natl. Acad. Sci. U.S.A.* 78:1444-45 (1981)); the regulatory sequences of the metallothionine gene (Brinster et al., *Nature* 296:39-42 (1982)); prokaryotic expression vectors such as the beta-lactamase promoter (Villa-Kamaroff et al., *Proc. Natl. Acad. Sci. U.S.A.*, 75:3727-31 (1978)); or the tac promoter (DeBoer et al., *Proc. Natl. Acad. Sci. U.S.A.*, 80:21-25 (1983)). Also of interest are the following animal transcriptional control regions, which exhibit tissue specificity and have been utilized in transgenic animals: the elastase I gene control region which is active in pancreatic acinar cells (Swift et al., *Cell* 38:639-46 (1984); Ornitz et al., *Cold Spring Harbor Symp. Quant. Biol.* 50:399-409 (1986); MacDonald, *Hepatology* 7:425-515 (1987)); the insulin gene control region which is active in pancreatic beta cells (Hanahan, *Nature* 315:115-22 (1985)); the immunoglobulin gene control region which is active in lymphoid cells (Grosschedl et al., *Cell* 38:647-58 (1984); Adames et al., *Nature* 318:533-38 (1985); Alexander et al., *Mol. Cell. Biol.*, 7:1436-44 (1987)); the mouse mammary tumor virus control region which is active in testicular, breast, lymphoid and mast cells (Leder et al., *Cell* 45:485-95 (1986)); the albumin gene control region which is active in liver (Pinkert et al., *Genes and Devel.* 1:268-76 (1987)); the alpha-feto-protein gene control region which is active in liver (Krumlauf et al., *Mol. Cell. Biol.*, 5:1639-48 (1985); Hammer et al., *Science* 235:53-58 (1987)); the alpha 1-antitrypsin gene control region which is active in the liver (Kelsey et al., *Genes and Devel.* 1:161-71, 1987)); the beta-globin gene control region which is active in myeloid cells (Mogam et al., *Nature* 315:338-40 (1985); Kollias et al., *Cell* 46:89-94 (1986)); the myelin basic protein gene control region which is active in oligodendrocyte cells in the brain (Readhead et al., *Cell* 48:703-12 (1987)); the myosin light chain-2 gene control region which is active in skeletal muscle (Sani,



- 34 -

*Nature* 314:283-86 (1985)); and the gonadotropic releasing hormone gene control region which is active in the hypothalamus (Mason et al., *Science* 234:1372-78 (1986)).

An enhancer sequence may be inserted into the vector to increase the transcription of a DNA encoding an FGF-like protein of the present invention by higher eukaryotes. Enhancers are cis-acting elements of DNA, usually about 10-300 bp in length, that act on the promoter to increase its transcription. Enhancers are relatively orientation and position independent. They have been found 5' and 3' to the transcription unit. Several enhancer sequences available from mammalian genes are known (e.g., globin, elastase, albumin, alpha-feto-protein and insulin). Typically, however, an enhancer from a virus will be used. The SV40 enhancer, the cytomegalovirus early promoter enhancer, the polyoma enhancer, and adenovirus enhancers are exemplary enhancing elements for the activation of eukaryotic promoters. While an enhancer may be spliced into the vector at a position 5' or 3' to FGF-like DNA, it is typically located at a site 5' from the promoter.

Expression vectors of the invention may be constructed from starting vectors such as a commercially available vector. Such vectors may or may not contain all of the desired flanking sequences. Where one or more of the flanking sequences set forth above are not already present in the vector to be used, they may be individually obtained and ligated into the vector. Methods used for obtaining each of the flanking sequences are well known to one skilled in the art.

Preferred vectors for practicing this invention are those which are compatible with bacterial, insect, and mammalian host cells. Such vectors include, *inter alia*, pCRII, pCR3, and pcDNA3.1 (Invitrogen, San Diego, CA), pBSII (Stratagene, La Jolla, CA), pET15 (Novagen, Madison, WI), pGEX (Pharmacia Biotech, Piscataway, NJ), pEGFP-N2 (Clontech, Palo Alto, CA), pETL (BlueBacII, Invitrogen), pDSR-alpha (PCT Pub. No. WO 90/14363) and pFastBacDual (Gibco-BRL, Grand Island, NY).

Additional possible vectors include, but are not limited to, cosmids,

- 35 -

plasmids, or modified viruses, but the vector system must be compatible with the selected host cell. Such vectors include, but are not limited to plasmids such as Bluescript<sup>®</sup> plasmid derivatives (a high copy number ColE1-based phagemid, Stratagene Cloning Systems, La Jolla CA), PCR cloning plasmids designed for  
5 cloning Taq-amplified PCR products (*e.g.*, TOPO<sup>™</sup> TA Cloning<sup>®</sup> Kit, PCR2.1<sup>®</sup> plasmid derivatives, Invitrogen, Carlsbad, CA), and mammalian, yeast or virus vectors such as a baculovirus expression system (pBacPAK plasmid derivatives, Clontech, Palo Alto, CA). The recombinant molecules can be introduced into host cells via transformation, transfection, infection, electroporation, or other known  
10 techniques.

After the vector has been constructed and a nucleic acid molecule encoding an FGF-like polypeptide has been inserted into the proper site of the vector, the completed vector may be inserted into a suitable host cell for amplification and/or polypeptide expression.

15 Host cells may be prokaryotic host cells (such as *E. coli*) or eukaryotic host cells (such as a yeast cell, an insect cell, or a vertebrate cell). The host cell, when cultured under appropriate conditions, synthesizes an FGF-like polypeptide which can subsequently be collected from the culture medium (if the host cell secretes it into the medium) or directly from the host cell producing it (if it is not  
20 secreted). Selection of an appropriate host cell will depend upon various factors, such as desired expression levels, polypeptide modifications that are desirable or necessary for activity, such as glycosylation or phosphorylation, and ease of folding into a biologically active molecule.

A number of suitable host cells are known in the art and many are  
25 available from the American Type Culture Collection (ATCC), Manassas, VA. Examples include mammalian cells, such as Chinese hamster ovary cells (CHO), CHO DHFR- cells (Urlaub et al., *Proc. Natl. Acad. Sci. U.S.A.* 97:4216-20 (1980)), human embryonic kidney (HEK) 293 or 293T cells, or 3T3 cells. The selection of suitable mammalian host cells and methods for transformation,  
30 culture, amplification, screening, product production and purification are known

- 36 -

in the art. Other suitable mammalian cell lines, are the monkey COS-1 and COS-7 cell lines, and the CV-1 cell line. Further exemplary mammalian host cells include primate cell lines and rodent cell lines, including transformed cell lines. Normal diploid cells, cell strains derived from *in vitro* culture of primary tissue, as well as primary explants, are also suitable. Candidate cells may be genotypically deficient in the selection gene, or may contain a dominantly acting selection gene. Other suitable mammalian cell lines include but are not limited to, mouse neuroblastoma N2A cells, HeLa, mouse L-929 cells, 3T3 lines derived from Swiss, Balb-c or NIH mice, BHK or HaK hamster cell lines. Each of these cell lines is known by and available to those skilled in the art of protein expression.

Similarly useful as host cells suitable for the present invention are bacterial cells. For example, the various strains of *E. coli* (e.g., HB101, DH5, DH10, and MC1061) are well known as host cells in the field of biotechnology. Various strains of *B. subtilis*, *Pseudomonas spp.*, other *Bacillus spp.*, *Streptomyces spp.*, and the like may also be employed in this method.

Many strains of yeast cells known to those skilled in the art are also available as host cells for expression of the polypeptides of the present invention. Preferred yeast cells include, for example, *Saccharomyces cerevisiae*.

Additionally, where desired, insect cell systems may be utilized in the methods of the present invention. Such systems are described, for example, in Kitts et al., *Biotechniques*, 14:810-17 (1993); Lucklow, *Curr. Opin. Biotechnol.* 4:564-72 (1993); and Lucklow et al., *J. Virol.*, 67:4566-79 (1993). Preferred insect cells are Sf-9 and Hi5 (Invitrogen).

Transformation or transfection of an expression vector for an FGF-like polypeptide into a selected host cell may be accomplished by well known methods including methods such as calcium chloride, electroporation, microinjection, lipofection or the DEAE-dextran method. The method selected will in part be a function of the type of host cell to be used. These methods and other suitable methods are well known to the skilled artisan, and are set forth, for example, in Sambrook et al., *supra*.

- 37 -

One may also use transgenic animals to express glycosylated FGF-like polypeptides. For example, one may use a transgenic milk-producing animal (a cow or goat, for example) and obtain the present glycosylated polypeptide in the animal milk. One may also use plants to produce FGF-like polypeptides, however, in general, the glycosylation occurring in plants is different from that produced in mammalian cells, and may result in a glycosylated product which is not suitable for human therapeutic use.

#### Polypeptide Production

Host cells comprising an FGF-like polypeptide expression vector (*i.e.*, transformed or transfected) may be cultured using standard media well known to the skilled artisan. The media will usually contain all nutrients necessary for the growth and survival of the cells. Suitable media for culturing *E. coli* cells are for example, Luria Broth (LB) and/or Terrific Broth (TB). Suitable media for culturing eukaryotic cells are RPMI 1640, MEM, DMEM, all of which may be supplemented with serum and/or growth factors as required by the particular cell line being cultured. A suitable medium for insect cultures is Grace's medium supplemented with yeastolate, lactalbumin hydrolysate, and/or fetal calf serum as necessary.

Typically, an antibiotic or other compound useful for selective growth of transfected or transformed cells is added as a supplement to the media. The compound to be used will be dictated by the selectable marker element present on the plasmid with which the host cell was transformed. For example, where the selectable marker element is kanamycin resistance, the compound added to the culture medium will be kanamycin. Other compounds for selective growth include ampicillin, tetracycline, and neomycin.

The amount of an FGF-like polypeptide produced by a host cell can be evaluated using standard methods known in the art. Such methods include, without limitation, Western blot analysis, SDS-polyacrylamide gel

- 38 -

electrophoresis, non-denaturing gel electrophoresis, HPLC separation, immunoprecipitation, and/or activity assays such as DNA binding gel shift assays.

If an FGF-like polypeptide has been designed to be secreted from the host cells, the majority of polypeptide may be found in the cell culture medium. If  
5 however, the FGF-like polypeptide is not secreted from the host cells, it will be present in the cytoplasm and/or the nucleus (for eukaryotic host cells) or in the cytosol (for gram-negative bacteria host cells).

For an FGF-like polypeptide situated in the host cell cytoplasm and/or nucleus, the host cells are typically first disrupted mechanically or with detergent  
10 to release the intra-cellular contents into a buffered solution. FGF-like polypeptide can then be isolated from this solution.

Purification of an FGF-like polypeptide from solution can be accomplished using a variety of techniques. If the polypeptide has been synthesized such that it contains a tag such as Hexahistidine (FGF-like  
15 polypeptide/hexaHis) or other small peptide such as FLAG (Eastman Kodak Co., New Haven, CT) or *myc* (Invitrogen) at either its carboxyl or amino terminus, it may essentially be purified in a one-step process by passing the solution through an affinity column where the column matrix has a high affinity for the tag or for the polypeptide directly (*i.e.*, a monoclonal antibody specifically recognizing  
20 FGF-like polypeptide). For example, polyhistidine binds with great affinity and specificity to nickel and thus an affinity column of nickel (such as the Qiagen® nickel columns) can be used for purification of FGF-like polypeptide/polyHis. *See, e.g., Current Protocols in Molecular Biology* § 10.11.8 (Ausubel et al., eds., John Wiley & Sons 1993).

25 Where an FGF-like polypeptide is prepared without a tag attached, and no antibodies are available, other well-known procedures for purification can be used. Such procedures include, without limitation, ion exchange chromatography, molecular sieve chromatography, HPLC, native gel electrophoresis in combination with gel elution, and preparative isoelectric focusing ("Isoprime")



- 39 -

machine/technique, Hoefer Scientific). In some cases, two or more of these techniques may be combined to achieve increased purity.

If an FGF-like polypeptide is produced intracellularly, the intracellular material (including inclusion bodies for gram-negative bacteria) can be extracted  
5 from the host cell using any standard technique known to the skilled artisan. For example, the host cells can be lysed to release the contents of the periplasm/cytoplasm by French press, homogenization, and/or sonication followed by centrifugation.

If an FGF-like polypeptide has formed inclusion bodies in the cytosol, the  
10 inclusion bodies can often bind to the inner and/or outer cellular membranes and thus will be found primarily in the pellet material after centrifugation. The pellet material can then be treated at pH extremes or with chaotropic agent such as a detergent, guanidine, guanidine derivatives, urea, or urea derivatives in the presence of a reducing agent such as dithiothreitol at alkaline pH or tris  
15 carboxyethyl phosphine at acid pH to release, break apart, and solubilize the inclusion bodies. The FGF-like polypeptide in its now soluble form can then be analyzed using gel electrophoresis, immunoprecipitation, or the like. If it is desired to isolate the FGF-like polypeptide, isolation may be accomplished using standard methods such as those set forth below and in Marston et al., *Meth. Enz.*,  
20 182:264-75 (1990).

In some cases, an FGF-like polypeptide may not be biologically active upon isolation. Various methods for "refolding" or converting the polypeptide to its tertiary structure and generating disulfide linkages can be used to restore biological activity. Such methods include exposing the solubilized polypeptide to  
25 a pH usually above 7 and in the presence of a particular concentration of a chaotrope. The selection of chaotrope is very similar to the choices used for inclusion body solubilization, but usually the chaotrope is used at a lower concentration and is not necessarily the same as chaotropes used for the solubilization. In most cases the refolding/oxidation solution will also contain a  
30 reducing agent or the reducing agent plus its oxidized form in a specific ratio to



- 40 -

generate a particular redox potential allowing for disulfide shuffling to occur in the formation of the protein's cysteine bridges. Some of the commonly used redox couples include cysteine/cystamine, glutathione (GSH)/dithiobis GSH, cupric chloride, dithiothreitol(DTT)/dithiane DTT, and 2-  
5 mercaptoethanol(bME)/dithio-b(ME). In many instances, a cosolvent may be used or may be needed to increase the efficiency of the refolding and the more common reagents used for this purpose include glycerol, polyethylene glycol of various molecular weights, arginine and the like.

If inclusion bodies are not formed to a significant degree upon expression  
10 of an FGF-like polypeptide, the polypeptide will be found primarily in the supernatant after centrifugation of the cell homogenate and may be further isolated from the supernatant using methods such as those set forth below.

In situations where it is preferable to partially or completely purify an FGF-like polypeptide such that it is partially or substantially free of contaminants,  
15 standard methods known to the one skilled in the art may be used. Such methods include, without limitation, separation by electrophoresis followed by electroelution, various types of chromatography (affinity, immunoaffinity, molecular sieve, and/or ion exchange), and/or high pressure liquid chromatography. In some cases, it may be preferable to use more than one of  
20 these methods for complete purification.

FGF-like polypeptides, fragments, and/or derivatives thereof may also be prepared by chemical synthesis methods (such as solid phase peptide synthesis) using techniques known in the art such as those set forth by Merrifield et al., *J. Am. Chem. Soc.* 85:2149 (1963); Houghten et al., *Proc Natl Acad. Sci. USA*  
25 82:5132 (1985); and Stewart and Young, *Solid Phase Peptide Synthesis* (Pierce Chemical Co. 1984). Such polypeptides may be synthesized with or without a methionine on the amino terminus. Chemically synthesized FGF-like polypeptides or fragments may be oxidized using methods set forth in these references to form disulfide bridges. FGF-like polypeptides, fragments or  
30 derivatives are expected to have comparable biological activity to the

- 41 -

corresponding FGF-like polypeptides, fragments or derivatives produced recombinantly or purified from natural sources, and thus may be used interchangeably with recombinant or natural FGF-like polypeptide.

Another means of obtaining FGF-like polypeptide is via purification from  
5 biological samples such as source tissues and/or fluids in which the FGF-like polypeptide is naturally found. Such purification can be conducted using methods for protein purification as described above. The presence of the FGF-like polypeptide during purification may be monitored using, for example, an antibody prepared against recombinantly produced FGF-like polypeptide or peptide  
10 fragments thereof.

#### Polypeptides

Polypeptides of the invention include isolated FGF-like polypeptides and polypeptides related thereto including fragments, variants, fusion polypeptides,  
15 and derivatives as defined hereinabove.

FGF-like polypeptide fragments of the invention may result from truncations at the amino terminus (with or without a leader sequence), truncations at the carboxyl terminus, and/or deletions internal to the polypeptide. In preferred embodiments, truncations and/or deletions comprise about 10 amino acids, or  
20 about 20 amino acid, or about 50 amino acids, or about 75 amino acids, or about 100 amino acids, or more than about 100 amino acids. The polypeptide fragments so produced will comprise about 25 contiguous amino acids, or about 50 amino acids, or about 75 amino acids, or about 100 amino acids, or about 150 amino acids, or about 200 amino acids. Such FGF-like polypeptide fragments may  
25 optionally comprise an amino terminal methionine residue.

FGF-like polypeptide variants of the invention include one or more amino acid substitutions, additions and/or deletions as compared to SEQ ID NO: 2 or SEQ ID NO: 4. In preferred embodiments, the variants have from 1 to 3, or from 1 to 5, or from 1 to 10, or from 1 to 15, or from 1 to 20, or from 1 to 25, or from 1  
30 to 50, or from 1 to 75, or from 1 to 100, or more than 100 amino acid

- 42 -

substitutions, insertions, additions and/or deletions, wherein the substitutions may be conservative, as defined above, non-conservative, or any combination thereof, and wherein the FGF-like polypeptide variant retains an FGF-like activity. The variants may have additions of amino acid residues either at the carboxyl terminus  
5 or at the amino terminus (with or without a leader sequence).

Preferred FGF-like polypeptide variants include glycosylation variants wherein the number and/or type of glycosylation sites has been altered compared to native FGF-like polypeptide. In one embodiment, FGF-like variants comprise a greater or a lesser number of N-linked glycosylation sites. An N-linked glycosylation site  
10 is characterized by the sequence: Asn-X-Ser or Thr, where the amino acid residue designated as "X" may be any type of amino acid except proline. Substitution(s) of amino acid residues to create this sequence provides a potential new site for addition of an N-linked carbohydrate chain. Alternatively, substitutions to eliminate this sequence will remove an existing N-linked carbohydrate chain.  
15 Also provided is a rearrangement of N-linked carbohydrate chains wherein one or more N-linked glycosylation sites (typically those that are naturally occurring) are eliminated and one or more new N-linked sites are created. Additional preferred FGF-like variants include cysteine variants, wherein one or more cysteine residues are deleted or substituted with another amino acid (e.g., serine). Cysteine variants  
20 are useful when FGF-like polypeptide must be refolded into a biologically active conformation such as after isolation of insoluble inclusive bodies. Cysteine variants generally have fewer cysteine residues than the native protein, and typically have an even number to minimize interactions resulting from unpaired cysteines.

25 One skilled in the art will be able to determine suitable variants of the native FGF-like polypeptide using well-known techniques. For example, one may be able to predict suitable areas of the molecule that may be changed without destroying biological activity. Also, one skilled in the art will realize that even areas that may be important for biological activity or for structure may be subject

- 43 -

to conservative amino acid substitutions without destroying the biological activity or without adversely affecting the polypeptide structure.

For predicting suitable areas of the molecule that may be changed without destroying activity, one skilled in the art may target areas not believed to be  
5 important for activity. For example, when similar polypeptides with similar activities from the same species or from other species are known, one skilled in the art may compare the amino acid sequence of FGF-like polypeptide to such similar polypeptides. After making such a comparison, one skilled in the art would be able to determine residues and portions of the molecules that are  
10 conserved among similar polypeptides. One skilled in the art would know that changes in areas of the FGF-like molecule that are not conserved would be less likely to adversely affect biological activity and/or structure. One skilled in the art would also know that, even in relatively conserved regions, one could have likely substituted chemically similar amino acids for the naturally occurring residues  
15 while retaining activity (conservative amino acid residue substitutions).

Also, one skilled in the art may review structure-function studies identifying residues in similar polypeptides that are important for activity or structure. In view of such a comparison, one skilled in the art can predict the importance of amino acid residues in FGF-like polypeptide that correspond to  
20 amino acid residues that are important for activity or structure in similar polypeptides. One skilled in the art may opt for chemically similar amino acid substitutions for such predicted important amino acid residues of FGF-like polypeptide.

If available, one skilled in the art can also analyze the three-dimensional  
25 structure and amino acid sequence in relation to that structure in similar polypeptides. In view of that information, one skilled in the art may be able to predict the alignment of amino acid residues of FGF-like polypeptide with respect to its three dimensional structure. One skilled in the art may choose not to make radical changes to amino acid residues predicted to be on the surface of the

- 44 -

protein, since such residues may be involved in important interactions with other molecules.

Moreover, one skilled in the art could generate test variants containing a single amino acid substitution at each amino acid residue. The variants could be screened using activity assays disclosed in this application. Such variants could be used to gather information about suitable variants. For example, if one discovered that a change to a particular amino acid residue resulted in destroyed activity, variants with such a change would be avoided. In other words, based on information gathered from such experiments, when attempting to find additional acceptable variants, one skilled in the art would have known the amino acids where further substitutions should be avoided either alone or in combination with other mutations.

FGF-like fusion polypeptides of the invention comprise FGF-like polypeptides, fragments, variants, or derivatives fused to a heterologous peptide or protein. Heterologous peptides and proteins include, but are not limited to: an epitope to allow for detection and/or isolation of an FGF-like fusion polypeptide; a transmembrane receptor protein or a portion thereof, such as an extracellular domain, or a transmembrane and intracellular domain; a ligand or a portion thereof which binds to a transmembrane receptor protein; an enzyme or portion thereof which is catalytically active; a protein or peptide which promotes oligomerization, such as leucine zipper domain; and a protein or peptide which increases stability, such as an immunoglobulin constant region. An FGF-like polypeptide may be fused to itself or to a fragment, variant, or derivative thereof. Fusions may be made either at the amino terminus or at the carboxyl terminus of an FGF-like polypeptide, and may be direct with no linker or adapter molecule or may be through a linker or adapter molecule, such as one or more amino acid residues up to about 20 amino acids residues, or up to about 50 amino acid residues. A linker or adapter molecule may also be designed with a cleavage site for a DNA restriction endonuclease or for a protease to allow for separation of the fused moieties.



- 45 -

In a further embodiment of the invention, an FGF-like polypeptide, fragment, variant and/or derivative is fused to an Fc region of human IgG. In one example, a human IgG hinge, CH2 and CH3 region may be fused at either the N-terminus or C-terminus of the FGF-like polypeptides using methods known to the skilled artisan. In another example, a portion of a hinge regions and CH2 and CH3 regions may be fused. The FGF-like Fc-fusion polypeptide so produced may be purified by use of a Protein A affinity column. In addition, peptides and proteins fused to an Fc region have been found to exhibit a substantially greater half-life *in vivo* than the unfused counterpart. Also, a fusion to an Fc region allows for dimerization/multimerization of the fusion polypeptide. The Fc region may be a naturally occurring Fc region, or may be altered to improve certain qualities, such as therapeutic qualities, circulation time, reduced aggregation, etc.

FGF-like polypeptide derivatives are included in the scope of the present invention. Such derivatives are chemically modified FGF-like polypeptide compositions in which FGF-like polypeptide is linked to a polymer. The polymer selected is typically water-soluble so that the protein to which it is attached does not precipitate in an aqueous environment, such as a physiological environment. The polymer may be of any molecular weight, and may be branched or unbranched. Included within the scope of FGF-like polypeptide polymers is a mixture of polymers. Preferably, for therapeutic use of the end-product preparation, the polymer will be pharmaceutically acceptable.

The water soluble polymer or mixture thereof may be selected from the group consisting of, for example, polyethylene glycol (PEG), monomethoxy-polyethylene glycol, dextran (such as low molecular weight dextran, of, for example about 6 kD), cellulose, or other carbohydrate based polymers, poly-(N-vinyl pyrrolidone) polyethylene glycol, propylene glycol homopolymers, a polypropylene oxide/ethylene oxide co-polymer, polyoxyethylated polyols (e.g., glycerol) and polyvinyl alcohol. Also encompassed by the invention are bifunctional PEG cross-linking molecules that may be used to prepare covalently attached FGF-like polypeptide multimers



For the acylation reactions, the polymer(s) selected should have a single reactive ester group. For reductive alkylation, the polymer(s) selected should have a single reactive aldehyde group. A reactive aldehyde is, for example, polyethylene glycol propionaldehyde, which is water stable, or mono C1-C10  
5 alkoxy or aryloxy derivatives thereof (*see* U.S. Patent No. 5,252,714).

Pegylation of FGF-like polypeptides may be carried out by any of the pegylation reactions known in the art, as described for example in the following references: Francis et al., *Focus on Growth Factors* 3, 4-10 (1992); EP 0 154 316; EP 0 401 384 and U.S. Patent No. 4,179,337. Pegylation may be carried out via  
10 an acylation reaction or an alkylation reaction with a reactive polyethylene glycol molecule (or an analogous reactive water-soluble polymer) as described below.

One water-soluble polymer for use herein is polyethylene glycol, abbreviated PEG. As used herein, polyethylene glycol is meant to encompass any of the forms of PEG that have been used to derivatize other proteins, such as  
15 mono-(C1-C10) alkoxy- or aryloxy-polyethylene glycol.

In general, chemical derivatization may be performed under any suitable conditions used to react a biologically active substance with an activated polymer molecule. Methods for preparing pegylated FGF-like polypeptides will generally comprise the steps of (a) reacting the polypeptide with polyethylene glycol (such  
20 as a reactive ester or aldehyde derivative of PEG) under conditions whereby FGF-like polypeptide becomes attached to one or more PEG groups, and (b) obtaining the reaction product(s). In general, the optimal reaction conditions for the acylation reactions will be determined based on known parameters and the desired result. For example, the larger the ratio of PEG: protein, the greater the  
25 percentage of poly-pegylated product.

In a preferred embodiment, the FGF-like polypeptide derivative will have a single PEG moiety at the amino terminus. *See* U.S. Patent No. 5,234,784, herein incorporated by reference.

Generally, conditions that may be alleviated or modulated by  
30 administration of the present FGF-like polypeptide derivative include those

described herein for FGF-like polypeptides. However, the FGF-like polypeptide derivative disclosed herein may have additional activities, enhanced or reduced biological activity, or other characteristics, such as increased or decreased half-life, as compared to the non-derivatized molecules.

5

### Antibodies

FGF-like polypeptides, fragments, variants, and derivatives may be used to prepare antibodies using methods known in the art. Thus, antibodies and antibody fragments that bind FGF-like polypeptides are within the scope of the present invention. Antibodies may be polyclonal, monospecific polyclonal, monoclonal,  
10 recombinant, chimeric, humanized, fully human, single chain and/or bispecific.

Polyclonal antibodies directed toward an FGF-like polypeptide generally are raised in animals (*e.g.*, rabbits or mice) by multiple subcutaneous or intraperitoneal injections of FGF-like polypeptide and an adjuvant. It may be  
15 useful to conjugate an FGF-like polypeptide, or a variant, fragment or derivative thereof to a carrier protein that is immunogenic in the species to be immunized, such as keyhole limpet hemocyanin, serum, albumin, bovine thyroglobulin, or soybean trypsin inhibitor. Also, aggregating agents such as alum are used to enhance the immune response. After immunization, the animals are bled and the  
20 serum is assayed for anti-FGF-like antibody titer.

Monoclonal antibodies directed toward FGF-like polypeptide are produced using any method that provides for the production of antibody molecules by continuous cell lines in culture. Examples of suitable methods for preparing monoclonal antibodies include hybridoma methods of Kohler, et al., *Nature*  
25 256:495-97 (1975), and the human B-cell hybridoma method, Kozbor, *J. Immunol.* 133:3001 (1984); Brodeur et al., *Monoclonal Antibody Production Techniques and Applications* 51-63 (Marcel Dekker 1987).

Also provided by the invention are hybridoma cell lines that produce monoclonal antibodies reactive with FGF-like polypeptides.

- 48 -

Monoclonal antibodies of the invention may be modified for use as therapeutics. One embodiment is a "chimeric" antibody in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequence in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequence in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (see U.S. Patent No. 4,816,567; Morrison, et al., *Proc. Natl. Acad. Sci. U.S.A.* 81: 6851-55 (1985)).

In another embodiment, a monoclonal antibody of the invention is a "humanized" antibody. Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source that is non-human. Humanization can be performed following methods known in the art (Jones, et al., *Nature* 321: 522-25 (1986); Riechmann, et al., *Nature* 332:323-27 (1988); Verhoeyen et al., *Science* 239:1534-36 (1988)), by substituting rodent complementarily-determining regions (CDRs) for the corresponding regions of a human antibody.

Also encompassed by the invention are fully human antibodies that bind FGF-like polypeptides, fragments, variants, and/or derivatives. Such antibodies are produced by immunization with an FGF-like antigen (optionally conjugated to a carrier) of transgenic animals (e.g., mice) that are capable of producing a repertoire of human antibodies in the absence of endogenous immunoglobulin production. See, e.g., Jakobovits, et al., *Proc. Natl. Acad. Sci. U.S.A.* 90: 2551-55 (1993); Jakobovits, et al., *Nature* 362:255-58 (1993); Bruggermann et al., *Year in Immuno.* 7:33 (1993). Human antibodies can also be produced in phage-display libraries (Hoogenboom et al., *J. Mol. Biol.* 227:381 (1991); Marks, et al., *J. Mol. Biol.* 222:581 (1991)).

Chimeric, CDR grafted and humanized antibodies are typically produced by recombinant methods. Nucleic acids encoding the antibodies are introduced

- 49 -

into host cells and expressed using materials and procedures described herein above. In a preferred embodiment, the antibodies are produced in mammalian host cells, such as CHO cells. Fully human antibodies may be produced by expression of recombinant DNA transfected into host cells or by expression in  
5 hybridoma cells as described above.

For diagnostic applications, in certain embodiments, anti-FGF-like antibodies typically will be labeled with a detectable moiety. The detectable moiety can be any one that is capable of producing, either directly or indirectly, a detectable signal. For example, the detectable moiety may be a radioisotope, such  
10 as  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ , or  $^{125}\text{I}$ , a fluorescent or chemiluminescent compound, such as fluorescein isothiocyanate, rhodamine, or luciferin; or an enzyme, such as alkaline phosphatase,  $\beta$ -galactosidase or horseradish peroxidase. Bayer, et al., *Meth. Enz.* 184: 138-63 (1990).

The anti-FGF-like antibodies of the invention may be employed in any  
15 known assay method, such as competitive binding assays, direct and indirect sandwich assays, and immunoprecipitation assays (Sola, *Monoclonal Antibodies: A Manual of Techniques* 147-58 (CRC Press 1987)) for detection and quantitation of FGF-like polypeptides. The antibodies will bind FGF-like polypeptides with an affinity that is appropriate for the assay method being employed.

20 Competitive binding assays rely on the ability of a labeled standard (e.g., an FGF-like polypeptide, or an immunologically reactive portion thereof) to compete with the test sample analyte (an FGF-like polypeptide) for binding with a limited amount of anti FGF-like antibody. The amount of an FGF-like polypeptide in the test sample is inversely proportional to the amount of standard  
25 that becomes bound to the antibodies. To facilitate determining the amount of standard that becomes bound, the antibodies typically are insolubilized before or after the competition, so that the standard and analyte that are bound to the antibodies may conveniently be separated from the standard and analyte which remain unbound.

- 50 -

Sandwich assays involve the use of two antibodies, each capable of binding to a different immunogenic portion, or epitope, of the protein to be detected and/or quantitated. In a sandwich assay, the test sample analyte is typically bound by a first antibody which is immobilized on a solid support, and thereafter a second antibody binds to the analyte, thus forming an insoluble three part complex. See, e.g., U.S. Patent No. 4,376,110. The second antibody may itself be labeled with a detectable moiety (direct sandwich assays) or may be measured using an anti-immunoglobulin antibody that is labeled with a detectable moiety (indirect sandwich assays). For example, one type of sandwich assay is an ELISA assay, in which case the detectable moiety is an enzyme.

The anti-FGF-like antibodies of the invention also are useful for *in vivo* imaging, wherein an antibody labeled with a detectable moiety is administered to an animal, preferably into the bloodstream, and the presence and location of the labeled antibody in the host is assayed. The antibody may be labeled with any moiety that is detectable in an animal, whether by nuclear magnetic resonance, radiology, or other detection means known in the art.

The invention also relates to a kit comprising anti-FGF-like antibodies and other reagents useful for detecting FGF-like polypeptide levels in biological samples. Such reagents may include a secondary activity, a detectable label, blocking serum, positive and negative control samples and detection reagents.

Antibodies of the invention may be used as therapeutics. Therapeutic antibodies are generally agonists or antagonists, in that they either enhance or reduce, respectively, at least one of the biological activities of an FGF-like polypeptide. In one embodiment, antagonist antibodies of the invention are antibodies or binding fragments thereof which are capable of specifically binding to an FGF-like polypeptide, fragment, variant, and/or derivative, and which are capable of inhibiting or eliminating the functional activity of an FGF-like polypeptide *in vivo* or *in vitro*. In preferred embodiments, an antagonist antibody will inhibit the functional activity of an FGF-like polypeptide at least about 50%, and preferably at least about 80%. In another embodiment antagonist antibodies



- 51 -

are capable of interacting with an FGF-like binding partner thereby inhibiting or eliminating FGF-like activity *in vitro* or *in vivo*. Agonist and antagonist anti-FGF-like antibodies are identified by screening assays described below.

5    Genetically Engineered Non-Human Animals

          Additionally included within the scope of the present invention are non-human animals such as mice, rats, or other rodents, rabbits, goats, sheep, or other farm animals, in which the genes encoding native FGF-like polypeptide have been disrupted (*i.e.*, "knocked out") such that the level of expression of FGF-like  
10   polypeptide is significantly decreased or completely abolished. Such animals may be prepared using techniques and methods such as those described in U.S. Patent No. 5,557,032.

          The present invention further includes non-human animals such as mice, rats, or other rodents, rabbits, goats, sheep, or other farm animals, in which a gene  
15   encoding a native form of FGF-like polypeptide for that animal or a heterologous FGF-like polypeptide gene is overexpressed by the animal, thereby creating a "transgenic" animal. Such transgenic animals may be prepared using well known methods such as those described in U.S. Patent No 5,489,743 and PCT Pub. No. WO 94/28122.

20    The present invention further includes non-human animals in which the promoter for one or more of the FGF-like polypeptides of the present invention is either activated or inactivated (*e.g.*, by using homologous recombination methods as described below) to alter the level of expression of one or more of the native FGF-like polypeptides.

25    Such non-human animals may be used for drug candidate screening. The impact of a drug candidate on the animal may be measured. For example, drug candidates may decrease or increase expression of the FGF-like polypeptide gene. In certain embodiments, the amount of FGF-like polypeptide or an FGF-like polypeptide fragment that is produced may be measured after exposure of the  
30   animal to the drug candidate. In certain embodiments, one may detect the actual



- 52 -

impact of the drug candidate on the animal. For example, overexpression of a particular gene may result in, or be associated with, a disease or pathological condition. In such cases, one may test a drug candidate's ability to decrease expression of the gene or its ability to prevent or inhibit a pathological condition.

5 In other examples, production of a particular metabolic product such as a fragment of a polypeptide, may result in, or be associated with, a disease or pathological condition. In such cases, one may test a drug candidate's ability to decrease production of such a metabolic product or its ability to prevent or inhibit a pathological condition.

10

#### Modulators of FGF-like polypeptide activity

In some situations, it may be desirable to identify molecules that are modulators, *i.e.*, agonists or antagonists, of the activity of FGF-like polypeptide.

Natural or synthetic molecules that modulate FGF-like polypeptide can be  
15 identified using one or more screening assays, such as those described below. Such molecules may be administered either in an *ex vivo* manner or in an *in vivo* manner by local or intravenous injection or by oral delivery, implantation device, or the like.

The following definition is used herein for describing the assays:

20 "Test molecule" refers to a molecule that is under evaluation for the ability to modulate (*i.e.*, increase or decrease) the activity of an FGF-like polypeptide. Most commonly, a test molecule will interact directly with an FGF-like polypeptide. However, it is also contemplated that a test molecule may also modulate FGF-like polypeptide activity indirectly, such as by affecting FGF-like  
25 gene expression, or by binding to an FGF-like binding partner (*e.g.*, receptor or ligand). In one embodiment, a test molecule will bind to an FGF-like polypeptide with an affinity constant of at least about  $10^{-6}$  M, preferably about  $10^{-8}$  M, more preferably about  $10^{-9}$  M, and even more preferably about  $10^{-10}$  M.

Methods for identifying compounds that interact with FGF-like  
30 polypeptides are encompassed by the invention. In certain embodiments, an FGF-

- 53 -

like polypeptide is incubated with a test molecule under conditions that permit interaction of the test molecule with an FGF-like polypeptide, and the extent of the interaction can be measured. The test molecule may be screened in a substantially purified form or in a crude mixture. Test molecules may be nucleic acid molecules, proteins, peptides, carbohydrates, lipids, or small molecular weight organic or inorganic compounds. Once a set of test molecules has been identified as interacting with an FGF-like polypeptide, the molecules may be further evaluated for their ability to increase or decrease FGF-like polypeptide activity.

10 Measurement of the interaction of test molecules with FGF-like polypeptides may be carried out in several formats, including cell-based binding assays, membrane binding assays, solution-phase assays and immunoassays. In general, test molecules are incubated with an FGF-like polypeptide for a specified period of time and FGF-like polypeptide activity is determined by one or more assays described herein for measuring biological activity.

Interaction of test molecules with FGF-like polypeptides may also be assayed directly using polyclonal or monoclonal antibodies in an immunoassay. Alternatively, modified forms of FGF-like polypeptides containing epitope tags as described above may be used in solution and immunoassays.

20 In certain embodiments, an FGF-like polypeptide agonist or antagonist may be a protein, peptide, carbohydrate, lipid, or small molecular weight molecule that interacts with FGF-like polypeptide to regulate its activity. Potential protein antagonists of FGF-like polypeptide include antibodies that interact with active regions of the polypeptide and inhibit or eliminate at least one activity of FGF-like polypeptide. Molecules which regulate FGF-like polypeptide expression may include nucleic acids which are complementary to nucleic acids encoding an FGF-like polypeptide, or are complementary to nucleic acid sequences which direct or control expression of FGF-like polypeptide, and which act as anti-sense regulators of expression.

In the event that FGF-like polypeptides display biological activity through interaction with a binding partner (*e.g.*, a receptor or a ligand), a variety of *in vitro* assays may be used to measure binding of an FGF-like polypeptide to a corresponding binding partner. These assays may be used to screen test molecules  
5 for their ability to increase or decrease the rate and/or the extent of binding of an FGF-like polypeptide to its binding partner. In one assay, an FGF-like polypeptide is immobilized by attachment to the bottom of the wells of a microtiter plate. Radiolabeled FGF-like binding partner (for example, iodinated FGF-like binding partner) and the test molecules can then be added either one at a  
10 time (in either order) or simultaneously to the wells. After incubation, the wells can be washed and counted using a scintillation counter for radioactivity to determine the extent of binding to FGF-like protein by its binding partner. Typically, the molecules will be tested over a range of concentrations and a series of control wells lacking one or more elements of the test assays can be used for  
15 accuracy in evaluation of the results. An alternative to this method involves reversing the "positions" of the proteins, *i.e.*, immobilizing FGF-like binding partner to the microtiter plate wells, incubating with the test molecule and radiolabeled FGF-like polypeptide, and determining the extent of FGF-like binding (*see, e.g., Current Protocols in Molecular Biology*, chap. 18 (Ausubel et  
20 al., eds., John Wiley & Sons 1995)).

As an alternative to radiolabeling, an FGF-like polypeptide or its binding partner may be conjugated to biotin and the presence of biotinylated protein can then be detected using streptavidin linked to an enzyme, such as horse radish peroxidase (HRP) or alkaline phosphatase (AP), that can be detected  
25 colorimetrically, or by fluorescent tagging of streptavidin. An antibody directed to an FGF-like polypeptide or to an FGF-like binding partner and that is conjugated to biotin may also be used and can be detected after incubation with enzyme-linked streptavidin linked to AP or HRP

An FGF-like polypeptide and an FGF-like binding partner may also be  
30 immobilized by attachment to agarose beads, acrylic beads, or other types of such

- 55 -

inert substrates. The substrate-protein complex can be placed in a solution containing the complementary protein and the test compound; after incubation, the beads can be precipitated by centrifugation, and the amount of binding between an FGF-like polypeptide and its binding partner can be assessed using the methods described above. Alternatively, the substrate-protein complex can be immobilized in a column and the test molecule and complementary protein passed over the column. Formation of a complex between an FGF-like polypeptide and its binding partner can then be assessed using any of the techniques set forth above, *i.e.*, radiolabeling, antibody binding, or the like.

Another *in vitro* assay that is useful for identifying a test molecule which increases or decreases formation of a complex between an FGF-like binding protein and an FGF-like binding partner is a surface plasmon resonance detector system such as the Biacore assay system (Pharmacia, Piscataway, NJ). The Biacore system may be carried out using the manufacturer's protocol. This assay essentially involves covalent binding of either FGF-like polypeptide or an FGF-like binding partner to a dextran-coated sensor chip that is located in a detector. The test compound and the other complementary protein can then be injected into the chamber containing the sensor chip either simultaneously or sequentially and the amount of complementary protein that binds can be assessed based on the change in molecular mass which is physically associated with the dextran-coated side of the sensor chip; the change in molecular mass can be measured by the detector system.

In some cases, it may be desirable to evaluate two or more test compounds together for their ability to increase or decrease formation of a complex between an FGF-like polypeptide and an FGF-like binding partner complex. In these cases, the assays set forth above can be readily modified by adding such additional test compounds either simultaneous with, or subsequent to, the first test compound. The remaining steps in the assay are as set forth above.

*In vitro* assays such as those described above may be used advantageously to screen rapidly large numbers of compounds for effects on complex formation

- 56 -

by FGF-like polypeptide and FGF-like binding partner. The assays may be automated to screen compounds generated in phage display, synthetic peptide, and chemical synthesis libraries.

Compounds which increase or decrease formation of a complex between  
5 an FGF-like polypeptide and an FGF-like binding partner may also be screened in cell culture using cells and cell lines expressing either FGF-like polypeptide or FGF-like binding partner. Cells and cell lines may be obtained from any mammal, but preferably will be from human or other primate, canine, or rodent sources. The binding of an FGF-like polypeptide to cells expressing FGF-like  
10 binding partner at the surface is evaluated in the presence or absence of test molecules and the extent of binding may be determined by, for example, flow cytometry using a biotinylated antibody to an FGF-like binding partner. Cell culture assays may be used advantageously to further evaluate compounds that score positive in protein binding assays described above.

15 Cell cultures can also be used to screen the impact of a drug candidate. For example, drug candidates may decrease or increase expression of the FGF-like polypeptide gene. In certain embodiments, the amount of FGF-like polypeptide or an FGF-like polypeptide fragment that is produced may be measured after exposure of the cell culture to the drug candidate. In certain embodiments, one  
20 may detect the actual impact of the drug candidate on the cell culture. For example, overexpression of a particular gene may have a particular impact on the cell culture. In such cases, one may test a drug candidate's ability to increase or decrease expression of the gene or its ability to prevent or inhibit a particular impact on the cell culture. In other examples, production of a particular metabolic  
25 product such as a fragment of a polypeptide, may result in, or be associated with, a disease or pathological condition. In such cases, one may test a drug candidate's ability to decrease production of such a metabolic product in a cell culture.



#### Cell Source Identification Using FGF-like Polypeptide

According to certain embodiments, it may be useful to be able to determine the source of a certain cell type. For example, it may be useful to  
5 determine the origin of a disease or pathological condition that may aid in selecting appropriate therapy. FGF-like polypeptide is specifically expressed in the liver (and weakly expressed in the lung). In certain embodiments, nucleic acid encoding an FGF-like polypeptide can be used as a probe to identify liver-derived cells by screening the nucleic acids of the cells with such a probe. In other  
10 embodiments, one may use the FGF-like polypeptide to make antibodies that are specific for FGF-like polypeptide. Such antibodies can be used to test for the presence of FGF-like polypeptide in cells, and thus, as a means for determining whether such cells were derived from the liver.

#### 15 FGF-like Polypeptide Compositions and Administration

Therapeutic compositions of FGF-like polypeptides are within the scope of the present invention. Such compositions may comprise a therapeutically effective amount of an FGF-like polypeptide, fragment, variant, or derivative in admixture with a pharmaceutically acceptable agent such as a pharmaceutically  
20 acceptable carrier. The carrier material may be water for injection, preferably supplemented with other materials common in solutions for administration to mammals. Typically, an FGF-like polypeptide therapeutic compound will be administered in the form of a composition comprising purified polypeptide, fragment, variant, or derivative in conjunction with one or more physiologically  
25 acceptable agents. Neutral buffered saline or saline mixed with serum albumin are exemplary appropriate carriers. Preferably, the product is formulated as a lyophilizate using appropriate excipients (e.g., sucrose). Other standard pharmaceutically acceptable agents such as carriers, diluents, and excipients may be included as desired. Other exemplary compositions comprise Tris buffer of



- 58 -

about pH 7.0-8.5, or acetate buffer of about pH 4.0-5.5, which may further include sorbitol or a suitable substitute therefor.

FGF-like polypeptide pharmaceutical compositions typically include a therapeutically or prophylactically effective amount of FGF-like protein in admixture with one or more pharmaceutically and physiologically acceptable formulation agents selected for suitability with the mode of administration. Suitable formulation materials or pharmaceutically acceptable agents include, but are not limited to, antioxidants, preservatives, coloring, flavoring and diluting agents, emulsifying agents, suspending agents, solvents, fillers, bulking agents, buffers, delivery vehicles, diluents, excipients and/or pharmaceutical adjuvants. For example, a suitable vehicle may be water for injection, physiological saline solution, or artificial cerebrospinal fluid, possibly supplemented with other materials common in compositions for parenteral administration. Neutral buffered saline or saline mixed with serum albumin are further exemplary vehicles. The term "pharmaceutically acceptable carrier" or "physiologically acceptable carrier" as used herein refers to a formulation agent(s) suitable for accomplishing or enhancing the delivery of the FGF-like protein as a pharmaceutical composition.

The primary solvent in a composition may be either aqueous or non-aqueous in nature. In addition, the vehicle may contain other formulation materials for modifying or maintaining the pH, osmolarity, viscosity, clarity, color, sterility, stability, rate of dissolution, or odor of the formulation. Similarly, the composition may contain additional formulation materials for modifying or maintaining the rate of release of FGF-like protein, or for promoting the absorption or penetration of FGF-like protein.

The FGF-like polypeptide compositions can be administered parentally. Alternatively, the compositions may be administered intravenously or subcutaneously. When systemically administered, the therapeutic compositions for use in this invention may be in the form of a pyrogen-free, parentally acceptable aqueous solution. The preparation of such pharmaceutically acceptable

- 59 -

protein solutions, with due regard to pH, isotonicity, stability and the like, is within the skill of the art.

Therapeutic formulations of FGF-like polypeptide compositions useful for practicing the present invention may be prepared for storage by mixing the  
5 selected composition having the desired degree of purity with optional physiologically acceptable carriers, excipients, or stabilizers (*Remington's Pharmaceutical Sciences* (18th Ed., A.R. Gennaro, ed., Mack Publishing Company 1990) in the form of a lyophilized cake or an aqueous solution. Acceptable carriers, excipients or stabilizers preferably are nontoxic to recipients  
10 and are preferably inert at the dosages and concentrations employed, and preferably include buffers such as phosphate, citrate, or other organic acids; antioxidants such as ascorbic acid; low molecular weight polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine,  
15 arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as Tween, pluronics or polyethylene glycol (PEG).

20 The optimal pharmaceutical formulation will be determined by one skilled in the art depending upon the intended route of administration, delivery format and desired dosage. See, e.g., *Remington's Pharmaceutical Sciences*, 1435-1712 (18th Ed., A.R. Gennaro, ed., Mack Publishing Company 1990). Such compositions may influence the physical state, stability, rate of *in vivo* release, and rate of *in*  
25 *vivo* clearance of the present FGF-like protein.

An effective amount of an FGF-like polypeptide composition to be employed therapeutically will depend, for example, upon the therapeutic objectives such as the indication for which the FGF-like polypeptide is being used, the route of administration, and the condition of the patient. Accordingly, it may  
30 be necessary for the therapist to titer the dosage and modify the route of

- 60 -

administration as required to obtain the optimal therapeutic effect. A typical dosage may range from about 0.1  $\mu\text{g/kg}$  to up to about 100 mg/kg or more, depending on the factors mentioned above. In other embodiments, the dosage may range from 1  $\mu\text{g/kg}$  up to about 100 mg/kg; or 5  $\mu\text{g/kg}$  up to about 100 mg/kg; or 0.1  $\mu\text{g/kg}$  up to about 100 mg/kg; or 1  $\mu\text{g/kg}$  up to about 100 mg/kg. Typically, a clinician will administer the composition until a dosage is reached that achieves the desired effect. The composition may therefore be administered as a single dose or as two or more doses (which may or may not contain the same amount of FGF-like polypeptide) over time, or as a continuous infusion via implantation device or catheter.

As further studies are conducted, information will emerge regarding appropriate dosage levels for treatment of various conditions in various patients, and the ordinary skilled worker, considering the therapeutic context, the type of disorder under treatment, the age and general health of the recipient, will be able to ascertain proper dosing.

The FGF-like polypeptide composition to be used for *in vivo* parenteral administration typically must be sterile. This is readily accomplished by filtration through sterile filtration membranes. Where the composition is lyophilized, sterilization using these methods may be conducted either prior to, or following, lyophilization and reconstitution. The composition for parenteral administration ordinarily will be stored in lyophilized form or in solution.

Therapeutic compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

Effective administration forms, such as (1) slow-release formulations, (2) inhalant mists, or (3) orally active formulations are also envisioned. The FGF-like polypeptide pharmaceutical composition also may be formulated for parenteral administration. Such parenterally administered therapeutic compositions are typically in the form of a pyrogen-free, parenterally acceptable aqueous solution comprising FGF-like polypeptide in a pharmaceutically acceptable vehicle. The

FGF-like polypeptide pharmaceutical compositions also may include particulate preparations of polymeric compounds such as polylactic acid, polyglycolic acid, etc., or the introduction of FGF-like polypeptide into liposomes. Hyaluronic acid may also be used, and this may have the effect of promoting sustained duration in the circulation.

A particularly suitable vehicle for parenteral injection is sterile distilled water in which FGF-like polypeptide is formulated as a sterile, isotonic solution, properly preserved. Yet another preparation may involve the formulation of FGF-like polypeptide with an agent, such as injectable microspheres, bio-erodible particles or beads, or liposomes, that provides for the controlled or sustained release of the protein product which may then be delivered as a depot injection. Other suitable means for the introduction of FGF-like polypeptide include implantable drug delivery devices that contain the FGF-like polypeptide.

The preparations of the present invention may include other components, for example parenterally acceptable preservatives, tonicity agents, cosolvents, wetting agents, complexing agents, buffering agents, antimicrobials, antioxidants and surfactants, as are well known in the art. For example, suitable tonicity enhancing agents include alkali metal halides (preferably sodium or potassium chloride), mannitol, sorbitol and the like. Suitable preservatives include, but are not limited to, benzalkonium chloride, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid and the like. Hydrogen peroxide may also be used as preservative. Suitable cosolvents are for example glycerin, propylene glycol, and polyethylene glycol. Suitable complexing agents are for example caffeine, polyvinylpyrrolidone, beta-cyclodextrin, or hydroxypropyl-beta-cyclodextrin. Suitable surfactants or wetting agents include sorbitan esters, polysorbates such as polysorbate 80, tromethamine, lecithin, cholesterol, tyloxapal and the like. The buffers can be conventional buffers such as borate, citrate, phosphate, bicarbonate, or Tris-HCl.

The formulation components are present in concentrations that are acceptable to the site of administration. For example, buffers are used to maintain

- 62 -

the composition at physiological pH or at slightly lower pH, typically within a pH range of from about 5 to about 8.

A pharmaceutical composition may be formulated for inhalation. For example, FGF-like polypeptide may be formulated as a dry powder for inhalation. 5 FGF-like polypeptide inhalation solutions may also be formulated in a liquefied propellant for aerosol delivery. In yet another formulation, solutions may be nebulized.

It is also contemplated that certain formulations containing FGF-like polypeptide may be administered orally. FGF-like polypeptide that is 10 administered in this fashion may be formulated with or without those carriers customarily used in the compounding of solid dosage forms such as tablets and capsules. For example, a capsule may be designed to release the active portion of the formulation at the point in the gastrointestinal tract when bioavailability is maximized and pre-systemic degradation is minimized. Additional agents may be 15 included to facilitate absorption of FGF-like polypeptide. Diluents, flavorings, low melting point waxes, vegetable oils, lubricants, suspending agents, tablet disintegrating agents, and binders may also be employed.

Another preparation may involve an effective quantity of FGF-like polypeptide in a mixture with non-toxic excipients that are suitable for the 20 manufacture of tablets. By dissolving the tablets in sterile water, or other appropriate vehicle, solutions can be prepared in unit dose form. Suitable excipients include, but are not limited to, inert diluents, such as calcium carbonate, sodium carbonate or bicarbonate, lactose, or calcium phosphate; or binding agents, such as starch, gelatin, or acacia; or lubricating agents such as 25 magnesium stearate, stearic acid, or talc.

Additional FGF-like polypeptide formulations will be evident to those skilled in the art, including formulations involving FGF-like polypeptide in combination with one or more other therapeutic agents. Techniques for formulating a variety of other sustained- or controlled-delivery means, such as 30 liposome carriers, bio-erodible microparticles or porous beads and depot



- 63 -

injections, are also known to those skilled in the art. See, for example, the Supersaxo et al. description of controlled release porous polymeric microparticles for the delivery of pharmaceutical compositions (See PCT Pub. No. WO 93/15722) the disclosure of which is hereby incorporated by reference.

5           Regardless of the manner of administration, the specific dose may be calculated according to body weight, body surface area, or organ size. Further refinement of the calculations necessary to determine the appropriate dosage for treatment involving each of the above mentioned formulations is routinely made by those of ordinary skill in the art and is within the ambit of tasks routinely  
10 performed by them. Appropriate dosages may be ascertained through use of appropriate dose-response data.

          The route of administration of the composition is in accord with known methods, for example, oral, injection or infusion by intravenous, intraperitoneal, intracerebral (intraparenchymal), intracerebroventricular, intramuscular,  
15 intraocular, intraarterial, or intralesional routes, or by sustained release systems or implantation device which may optionally involve the use of a catheter. Where desired, the compositions may be administered continuously by infusion, bolus injection or by implantation device.

          One may further administer the present pharmaceutical compositions by  
20 pulmonary administration, *see, e.g.*, PCT Pub. WO 94/20069, which discloses pulmonary delivery of chemically modified proteins, herein incorporated by reference. For pulmonary delivery, the particle size should be suitable for delivery to the distal lung. For example, the particle size may be from 1  $\mu\text{m}$  to 5  $\mu\text{m}$ . However, larger particles may be used, for example, if each particle is fairly  
25 porous.

          Alternatively or additionally, the composition may be administered locally via implantation into the affected area of a membrane, sponge, or other appropriate material on to which FGF-like polypeptide has been absorbed or encapsulated.



- 64 -

Where an implantation device is used, the device may be implanted into any suitable tissue or organ, and delivery of FGF-like polypeptide may be directly through the device via bolus, or via continuous administration, or via catheter using continuous infusion.

5 FGF-like polypeptide may be administered in a sustained release formulation or preparation. Suitable examples of sustained-release preparations include semipermeable polymer matrices in the form of shaped articles, for example, films, or microcapsules. Sustained release matrices include polyesters, hydrogels, polylactides (U.S. Patent No. 3,773,919, EP Patent No. 58,481),  
10 copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman et al, *Biopolymers* 22: 547-56 (1983)), poly (2-hydroxyethyl-methacrylate) (Langer et al., *J. Biomed. Mater. Res.* 15: 167-277 (1981) and Langer, *Chem. Tech.* 12: 98-105 (1982)), ethylene vinyl acetate (Langer et al., *supra*) or poly-D(-)-3-hydroxybutyric acid (EP Patent No. 133,988). Sustained-release compositions  
15 also may include liposomes, which can be prepared by any of several methods known in the art (*see, e.g.*, Epstein et al., *Proc. Natl. Acad. Sci. U.S.A.* 82:3688-92 (1985); EP Patent Nos. 36,676; 88,046; and 143,949).

The FGF-like polypeptides, fragments thereof, variants, and derivatives, may be employed alone, together, or in combination with other pharmaceutical  
20 compositions. The FGF-like polypeptides, fragments, variants, and derivatives may be used in combination with cytokines, growth factors, antibiotics, anti-inflammatories, and/or chemotherapeutic agents as is appropriate for the indication being treated.

In some cases, it may be desirable to use FGF-like polypeptide  
25 compositions in an *ex vivo* manner. Here, cells, tissues, or organs that have been removed from the patient are exposed to FGF-like polypeptide compositions after which the cells, tissues and/or organs are subsequently implanted back into the patient.

In other cases, an FGF-like polypeptide may be delivered through  
30 implanting into patients certain cells that have been genetically engineered, using

- 65 -

methods such as those described herein, to express and secrete the polypeptides, fragments, variants, or derivatives. Such cells may be animal or human cells, and may be derived from the patient's own tissue or from another source, either human or non-human. Optionally, the cells may be immortalized. However, in order to  
5 decrease the chance of an immunological response, it is preferred that the cells be encapsulated to avoid infiltration of surrounding tissues. The encapsulation materials are typically biocompatible, semi-permeable polymeric enclosures or membranes that allow release of the protein product(s) but prevent destruction of the cells by the patient's immune system or by other detrimental factors from the  
10 surrounding tissues.

Methods used for membrane encapsulation of cells are familiar to the skilled artisan, and preparation of encapsulated cells and their implantation in patients may be accomplished without undue experimentation. *See, e.g.,* U.S. Patent Nos. 4,892,538; 5,011,472; and 5,106,627. A system for encapsulating  
15 living cells is described in PCT Pub. No. WO 91/10425 (Aebischer et al.). Techniques for formulating a variety of other sustained or controlled delivery means, such as liposome carriers, bio-erodible particles or beads, are also known to those in the art, and are described. The cells, with or without encapsulation, may be implanted into suitable body tissues or organs of the patient.

20 As discussed above, it may be desirable to treat isolated cell populations such as stem cells, leukocytes, red blood cells, bone marrow, chondrocytes, neurons, pancreatic islets, liver cells and the like with one or more FGF-like polypeptides, variants, derivatives and/or fragments. This can be accomplished by exposing the isolated cells to the polypeptide, variant, derivative, or fragment  
25 directly, where it is in a form that is permeable to or acts upon the cell membrane.

Additional objects of the present invention relate to methods for both the *in vitro* production of therapeutic proteins by means of homologous recombination and for the production and delivery of therapeutic proteins by gene therapy.

It is further envisioned that FGF-like protein may be produced by  
30 homologous recombination, or with recombinant production methods utilizing

- 66 -

control elements introduced into cells already containing DNA encoding FGF-like polypeptide. For example, homologous recombination methods may be used to modify a cell that contains a normally transcriptionally silent FGF-like gene, or under expressed gene, and thereby produce a cell that expresses therapeutically efficacious amounts of FGF-like polypeptide. Homologous recombination is a technique originally developed for targeting genes to induce or correct mutations in transcriptionally active genes (Kucherlapati, *Prog. in Nucl. Acid Res. and Mol. Biol.* 36:301 (1989)). The basic technique was developed as a method for introducing specific mutations into specific regions of the mammalian genome (Thomas et al., *Cell* 44:419-28 (1986); Thomas and Capecchi, *Cell* 51:503-12, (1987); Doetschman et al., *Proc. Natl. Acad. Sci. U.S.A.* 85:8583-87 (1988)) or to correct specific mutations within defective genes (Doetschman et al., *Nature* 330:576-78 (1987)). Exemplary homologous recombination techniques are described in U.S. Patent No. 5,272,071 (EP Patent No. 91 90 3051, EP Publication No. 505 500; PCT/US90/07642, PCT Pub. No. WO 91/09955).

Through homologous recombination, the DNA sequence to be inserted into the genome can be directed to a specific region of the gene of interest by attaching it to targeting DNA. The targeting DNA is a nucleotide sequence that is complementary (homologous) to a region of the genomic DNA. Small pieces of targeting DNA that are complementary to a specific region of the genome are put in contact with the parental strand during the DNA replication process. It is a general property of DNA that has been inserted into a cell to hybridize, and therefore, recombine with other pieces of endogenous DNA through shared homologous regions. If this complementary strand is attached to an oligonucleotide that contains a mutation or a different sequence or an additional nucleotide, it too is incorporated into the newly synthesized strand as a result of the recombination. As a result of the proofreading function, it is possible for the new sequence of DNA to serve as the template. Thus, the transferred DNA is incorporated into the genome.

Attached to these pieces of targeting DNA are regions of DNA that may

- 67 -

interact with the expression of a FGF-like protein. For example, a promoter/enhancer element, a suppresser, or an exogenous transcription modulatory element is inserted in the genome of the intended host cell in proximity and orientation sufficient to influence the transcription of DNA  
5 encoding the desired FGF-like protein. The control element controls a portion of the DNA present in the host cell genome. Thus, the expression of FGF-like protein may be achieved not by transfection of DNA that encodes the FGF-like gene itself, but rather by the use of targeting DNA (containing regions of homology with the endogenous gene of interest) coupled with DNA regulatory  
10 segments that provide the endogenous gene sequence with recognizable signals for transcription of a FGF-like protein.

In an exemplary method, expression of a desired targeted gene in a cell (*i.e.*, a desired endogenous cellular gene) is altered by the introduction, by homologous recombination into the cellular genome at a preselected site, of DNA  
15 which includes at least a regulatory sequence, an exon and a splice donor site. These components are introduced into the chromosomal (genomic) DNA in such a manner that this, in effect, results in production of a new transcription unit (in which the regulatory sequence, the exon, and the splice donor site present in the DNA construct are operatively linked to the endogenous gene). As a result of  
20 introduction of these components into the chromosomal DNA, the expression of the desired endogenous gene is altered.

Altered gene expression, as used herein, encompasses activating (or causing to be expressed) a gene which is normally silent (unexpressed) in the cell as obtained, increasing expression of a gene which may include expressing a gene  
25 that is not expressed at physiologically significant levels in the cell as obtained, changing the pattern of regulation or induction such that it is different than occurs in the cell as obtained, and reducing (including eliminating) expression of a gene which is expressed in the cell as obtained.

The present invention further relates to DNA constructs useful in the  
30 method of altering expression of a target gene. In certain embodiments, the

- 68 -

exemplary DNA constructs comprise: (a) a targeting sequence, (b) a regulatory sequence, (c) an exon, and (d) an unpaired splice-donor site. The targeting sequence in the DNA construct directs the integration of elements (a)-(d) into a target gene in a cell such that the elements (b)-(d) are operatively linked to sequences of the endogenous target gene. In another embodiment, the DNA constructs comprise: (a) a targeting sequence, (b) a regulatory sequence, (c) an exon, (d) a splice-donor site, (e) an intron, and (f) a splice-acceptor site, wherein the targeting sequence directs the integration of elements (a)-(f) such that the elements of (b)-(f) are operatively linked to the endogenous gene. The targeting sequence is homologous to the preselected site in the cellular chromosomal DNA with which homologous recombination is to occur. In the construct, the exon is generally 3' of the regulatory sequence and the splice-donor site is 3' of the exon.

If the sequence of a particular gene is known, such as the nucleic acid sequence of FGF-like polypeptide presented herein, a piece of DNA that is complementary to a selected region of the gene can be synthesized or otherwise obtained, such as by appropriate restriction of the native DNA at specific recognition sites bounding the region of interest. This piece serves as a targeting sequence upon insertion into the cell and will hybridize to its homologous region within the genome. If this hybridization occurs during DNA replication, this piece of DNA, and any additional sequence attached thereto, will act as an Okazaki fragment and will be backstitched into the newly synthesized daughter strand of DNA. The present invention, therefore, includes nucleotides encoding a FGF-like molecule, which nucleotides may be used as targeting sequences.

FGF-like polypeptide cell therapy – for example, implantation of cells producing FGF-like polypeptide – is also contemplated. This embodiment would involve implanting into the cells of a patient a construct capable of synthesizing and secreting a biologically active form of FGF-like polypeptide. Such FGF-like polypeptide-producing cells may be cells that are natural producers of FGF-like polypeptide or may be recombinant cells whose ability to produce FGF-like polypeptide has been augmented by transformation with a gene encoding the



- 69 -

desired FGF-like molecule or with a gene augmenting the expression of FGF-like polypeptide. Such a modification may be accomplished by means of a vector suitable for delivering the gene as well as promoting its expression and secretion. In order to minimize a potential immunological reaction in a patient that is being administered an FGF-like protein or polypeptide of a foreign species, it is preferred that the natural cells producing FGF-like polypeptide be of human origin and produce human FGF-like polypeptide. Likewise, it is preferred that the recombinant cells producing FGF-like polypeptide are transformed with an expression vector containing a gene encoding a human FGF-like molecule.

10           Implanted cells may be encapsulated to avoid infiltration of surrounding tissue. Human or non-human animal cells may be implanted in patients in biocompatible, semipermeable polymeric enclosures or membranes that allow release of FGF-like polypeptide, but that prevent destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding  
15 tissue. Alternatively, the patient's own cells, transformed to produce FGF-like polypeptide *ex vivo*, could be implanted directly into the patient without such encapsulation.

Techniques for the encapsulation of living cells are known in the art, and the preparation of the encapsulated cells and their implantation in a patient may be  
20 accomplished without undue experimentation. For example, Baetge et al. (PCT Pub. No. WO 95/05452, the disclosure of which is hereby incorporated by reference) describe membrane capsules containing genetically engineered cells for the effective delivery of biologically active molecules. The capsules are biocompatible and are easily retrievable. The capsules encapsulate cells  
25 transfected with recombinant DNA molecules comprising DNA sequences coding for biologically active molecules operatively linked to promoters that are not subject to down regulation *in vivo* upon implantation into a mammalian host. The devices provide for the delivery of the molecules from living cells to specific sites within a recipient. In addition, see U.S. Patent Nos. 4,892,538; 5,011,472; and  
30 5,106,627. A system for encapsulating living cells is described in PCT Pub. No.

- 70 -

WO 91/10425 (Aebischer et al.). See also, PCT Pub. No. WO 91/10470 (Aebischer et al.); Winn et al., *Exper. Neurol.* 113:322-29 (1991); Aebischer et al., *Exper. Neurol.*, 111:269-75 (1991); and Tresco et al., *ASAIO* 38:17-23 (1992).

*In vivo* and *in vitro* gene therapy delivery of FGF-like polypeptide is also envisioned. *In vivo* gene therapy may be accomplished by introducing the gene encoding FGF-like polypeptide into cells via local injection of a polynucleotide molecule or other appropriate delivery vectors (Hefti, *J. Neurobiology* 25:1418-35 (1994)). For example, a polynucleotide molecule encoding FGF-like protein may be contained in an adeno-associated virus vector for delivery to the targeted cells (see, e.g., Johnson, PCT Pub. No. WO 95/34670; PCT App. No. PCT/US95/07178). The recombinant adeno-associated virus (AAV) genome typically contains AAV inverted terminal repeats flanking a DNA sequence encoding FGF-like polypeptide operably linked to functional promoter and polyadenylation sequences.

Alternative viral vectors include, but are not limited to, retrovirus, adenovirus, herpes simplex virus, and papilloma virus vectors. U.S. Patent No. 5,672,344 describes an *in vivo* viral-mediated gene transfer system involving a recombinant neurotrophic HSV-1 vector. U.S. Patent No. 5,399,346 provides examples of a process for providing a patient with a therapeutic protein by the delivery of human cells which have been treated *in vitro* to insert a DNA segment encoding a therapeutic protein. Additional methods and materials for the practice of gene therapy techniques are described in U.S. Patent No. 5,631,236 (involving adenoviral vectors); U.S. Patent No. 5,672,510 (involving retroviral vectors); and U.S. Patent 5,635,399 (involving retroviral vectors expressing cytokines).

Nonviral delivery methods include liposome-mediated transfer, naked DNA delivery (direct injection), receptor-mediated transfer (ligand-DNA complex), electroporation, calcium phosphate precipitation, and microparticle bombardment (e.g., gene gun). Gene therapy materials and methods may also include inducible promoters, tissue-specific enhancer-promoters, DNA sequences designed for site-specific integration, DNA sequences capable of providing a

selective advantage over the parent cell, labels to identify transformed cells, negative selection systems and expression control systems (safety measures), cell-specific binding agents (for cell targeting), cell-specific internalization factors, transcription factors to enhance expression by a vector as well as methods of  
5 vector manufacture. Such additional methods and materials for the practice of gene therapy techniques are described in U.S. Patent No. 4,970,154 (electroporation techniques); PCT Pub. No. WO 96/40958 (nuclear ligands); U.S. Patent No. 5,679,559 (concerning a lipoprotein-containing system for gene delivery); U.S. Patent No. 5,676,954 (involving liposome carriers); U.S. Patent  
10 No. 5,593,875 (concerning methods for calcium phosphate transfection); and U.S. Patent No. 4,945,050 (wherein biologically active particles are propelled at cells at a speed whereby the particles penetrate the surface of the cells and become incorporated into the interior of the cells). Expression control techniques include chemical induced regulation (*see, e.g.*, PCT Pub. Nos. WO 96/41865 and WO  
15 97/31899), the use of a progesterone antagonist in a modified steroid hormone receptor system (*see, e.g.*, U.S. Patent No. 5,364,791), ecdysone control systems (*see, e.g.*, PCT Pub. No. WO 96/37609), and positive tetracycline-controllable transactivators (*see, e.g.*, U.S. Patent No. 5,589,362; U.S. Patent No. 5,650,298; and U.S. Patent No. 5,654,168).

20 It is also contemplated that FGF-like polypeptide gene therapy or cell therapy can further include the delivery of a second protein. For example, the host cell may be modified to express and release both FGF-like polypeptide and a second protein, for example insulin-like growth factor 1 (IGF-1). Alternatively, both FGF-like polypeptide and a second protein may be expressed in and released  
25 from separate cells. Such cells may be separately introduced into the patient or the cells may be contained in a single implantable device, such as the encapsulating membrane described above.

One manner in which gene therapy can be applied is to use the FGF-like gene (either genomic DNA, cDNA, and/or synthetic DNA encoding a FGF-like  
30 polypeptide, or a fragment, variant, or derivative thereof) which may be operably

- 72 -

linked to a constitutive or inducible promoter to form a "gene therapy DNA construct." The promoter may be homologous or heterologous to the endogenous FGF-like gene, provided that it is active in the cell or tissue type into which the construct will be inserted. Other components of the gene therapy DNA construct may optionally include: DNA molecules designed for site-specific integration (e.g., endogenous flanking sequences useful for homologous recombination), a tissue-specific promoter, enhancers, silencers, DNA molecules capable of providing a selective advantage over the parent cell, DNA molecules useful as labels to identify transformed cells, negative selection systems, cell specific binding agents (as, for example, for cell targeting), cell-specific internalization factors, and transcription factors to enhance expression by a vector as well as factors to enable vector manufacture.

This gene therapy DNA construct can then be introduced into the patient's cells (either *ex vivo* or *in vivo*). One means for introducing the gene therapy DNA construct is via viral vectors. Suitable viral vectors typically used in gene therapy for delivery of gene therapy DNA constructs include, without limitation, adenovirus, adeno-associated virus, herpes simplex virus, lentivirus, papilloma virus, and retrovirus vectors. Some of these vectors, such as retroviral vectors, will deliver the gene therapy DNA construct to the chromosomal DNA of the patient's cells, and the gene therapy DNA construct can integrate into the chromosomal DNA; other vectors will function as episomes and the gene therapy DNA construct will remain in the cytoplasm.

Another means to increase endogenous FGF-like polypeptide expression in a cell via gene therapy is to insert one or more enhancer elements into the FGF-like polypeptide promoter, where the enhancer elements can serve to increase transcriptional activity of the FGF-like polypeptide gene. The enhancer elements used will be selected based on the tissue in which one desires to activate the gene – enhancer elements known to confer promoter activation in that tissue will be selected. For example, if a gene encoding an FGF-like polypeptide is to be "turned on" in T-cells, the *lck* promoter enhancer element may be used. Here, the

- 73 -

functional portion of the transcriptional element to be added may be inserted into a fragment of DNA containing the FGF-like polypeptide promoter (and optionally, inserted into a vector and/or 5' and/or 3' flanking sequences, etc.) using standard cloning techniques. This construct, known as a "homologous recombination  
5 construct," can then be introduced into the desired cells either *ex vivo* or *in vivo*.

Gene therapy can be used to decrease FGF-like polypeptide expression by modifying the nucleotide sequence of the endogenous promoter. Such modification is typically accomplished via homologous recombination methods. For example, a DNA molecule containing all or a portion of the promoter of the  
10 FGF-like gene selected for inactivation can be engineered to remove and/or replace pieces of the promoter that regulate transcription. Here, for example, the TATA box and/or the binding site of a transcriptional activator of the promoter may be deleted using standard molecular biology techniques; such deletion can inhibit promoter activity thereby repressing transcription of the corresponding  
15 FGF-like gene. Deletion of the TATA box or transcription activator binding site in the promoter may be accomplished by generating a DNA construct comprising all or the relevant portion of the FGF-like polypeptide promoter (from the same or a related species as the FGF-like gene to be regulated) in which one or more of the TATA box and/or transcriptional activator binding site nucleotides are mutated  
20 via substitution, deletion and/or insertion of one or more nucleotides such that the TATA box and/or activator binding site has decreased activity or is rendered completely inactive. This construct, which also will typically contain at least about 500 bases of DNA that correspond to the native (endogenous) 5' and 3' DNA sequences adjacent to the promoter segment that has been modified, may be  
25 introduced into the appropriate cells (either *ex vivo* or *in vivo*) either directly or via a viral vector as described above. Typically, integration of the construct into the genomic DNA of the cells will be via homologous recombination, where the 5' and 3' DNA sequences in the promoter construct can serve to help integrate the modified promoter region via hybridization to the endogenous chromosomal  
30 DNA.



- 74 -

Other gene therapy methods may also be employed where it is desirable to inhibit the activity of one or more FGF-like polypeptides. For example, antisense DNA or RNA molecules, which have a sequence that is complementary to at least a portion of the selected FGF-like polypeptide gene can be introduced into the cell. Typically, each such antisense molecule will be complementary to the start site (5' end) of each selected FGF-like gene. When the antisense molecule then hybridizes to the corresponding FGF-like mRNA, translation of this mRNA is prevented.

Alternatively, gene therapy may be employed to create a dominant-negative inhibitor of one or more FGF-like polypeptides. In this situation, the DNA encoding a mutant full length or truncated polypeptide of each selected FGF-like polypeptide can be prepared and introduced into the cells of a patient using either viral or non-viral methods as described above. Each such mutant is typically designed to compete with endogenous polypeptide in its biological role.

15

#### Uses of FGF-like Nucleic Acids and Polypeptides

Nucleic acid molecules of the invention may be used to map the locations of the FGF-like gene and related genes on chromosomes. Mapping may be done by techniques known in the art, such as PCR amplification and *in situ* hybridization.

20

The nucleic acid molecules are also used as anti-sense inhibitors of FGF-like polypeptide expression. Such inhibition may be effected by nucleic acid molecules that are complementary to and hybridize to expression control sequences (triple helix formation) or to FGF-like mRNA. Anti-sense probes may be designed by available techniques using the sequence of the FGF-like genes disclosed herein. Anti-sense inhibitors provide information relating to the decrease or absence of an FGF-like polypeptide in a cell or organism.

25

Hybridization probes may be prepared using an FGF-like gene sequence as provided herein to screen cDNA, genomic or synthetic DNA libraries for related sequences. Regions of the DNA and/or amino acid sequence of FGF-like

30

- 75 -

polypeptide that exhibit significant identity to known sequences are readily determined using sequence alignment algorithms disclosed above and those regions may be used to design probes for screening.

The nucleic acid molecules of the invention may be used for gene therapy.

- 5 Nucleic acid molecules that express FGF-like polypeptide *in vivo* provide information relating to the effects of the polypeptide in cells or organisms.

FGF-like nucleic acid molecules, fragments, variants, and/or derivatives that do not themselves encode biologically active polypeptides may be useful as hybridization probes in diagnostic assays to test, either qualitatively or  
10 quantitatively, for the presence of FGF-like DNA or corresponding RNA in mammalian tissue or bodily fluid samples.

FGF-like polypeptides, fragments, variants, and/or derivatives may be used to prevent or treat cirrhosis or other toxic insult of the liver; inflammatory bowel disease, mucositis, Crohn's disease, or other gastrointestinal abnormality;  
15 diabetes; obesity; neurodegenerative diseases; wounds; damage to the corneal epithelium, lens, or retinal tissue; damage to renal tubules as a result of acute tubular necrosis; hematopoietic cell reconstitution following chemotherapy; wasting syndromes (for example, cancer associated cachexia), multiple sclerosis, myopathies; short stature, delayed maturation, excessive growth (for example,  
20 acromegaly), premature maturation; alopecia; diseases or abnormalities of androgen target organs; infantile respiratory distress syndrome, bronchopulmonary dysplasia, acute respiratory distress syndrome, or other lung abnormalities; tumors of the eye or other tissues; atherosclerosis; hypercholesterolemia; diabetes; obesity; stroke; osteoporosis; osteoarthritis; degenerative joint disease; muscle  
25 atrophy; sarcopenia; decreased lean body mass; baldness; wrinkles; increased fatigue; decreased stamina; decreased cardiac function; immune system dysfunction; cancer; Parkinson's disease; senile dementia; Alzheimer's disease; and decreased cognitive function.

FGF-like polypeptide fragments, variants, and/or derivatives, whether  
30 biologically active or not, are useful for preparing antibodies that bind to an FGF-

- 76 -

like polypeptide. The antibodies may be used for *in vivo* and *in vitro* diagnostic purposes, including, but not limited to, use in labeled form to detect the presence of FGF-like polypeptide in a body fluid or cell sample. The antibodies may also be used to prevent or treat conditions that may be associated with an increase in FGF-like polypeptide expression or activity. The antibodies may bind to an FGF-like polypeptide so as to diminish or block at least one activity characteristic of an FGF-like polypeptide, or may bind to a polypeptide to increase an activity.

A deposit of cDNA encoding FGF-like polypeptide has been made with the American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209 on September 3, 1999 and having accession No. PTA-626.

The following examples are intended for illustration purposes only, and should not be construed as limiting the scope of the invention in any way.

#### Example 1: Cloning of the Murine FGF-like Polypeptide Gene

Generally, materials and methods as described in Sambrook et al. *supra* were used to clone and analyze the gene encoding rat FGF-like polypeptide.

Sequences encoding the murine FGF-like polypeptide were isolated from a mouse regenerating liver cDNA library by screening the library in a kFGF signal trap system (U.S. Patent App. No. 09/026,958). This primary screening technique enriched for clones encoding signal peptide-containing secreted proteins.

A primary library (Tmr11) was constructed in the kFGF vector as follows. Regenerating mouse liver was removed 24 hours after partial hepatectomy, and poly A<sup>+</sup> RNA prepared using a commercially available RNA extraction kit and mRNA purification kit (Pharmacia Biotech). A cDNA library was prepared using the SuperScript<sup>TM</sup> Plasmid System for cDNA Synthesis and Plasmid Cloning (Gibco BRL) with some modifications. First strand reactions were performed using 3 µg of poly A<sup>+</sup> RNA and 500 µg of the primer 5'-G-G-A-A-G-G-A-A-A-A-A-A-G-C-G-G-C-C-G-C-A-A-C-A-N-N-N-N-N-N-N-N-N-N-3' (SEQ ID NO: 34). Following second strand synthesis, the cDNA was ligated to a Sal I adapter, digested with Not I, and then size fractionated by agarose gel electrophoresis.

- 77 -

Fractionated cDNA, ranging from 0.2 to 0.8 kb in size, was purified using a Qiagen gel extraction kit and then ligated to the kFGF vector as follows. In a 20  $\mu$ l reaction, 50  $\mu$ g of vector DNA (previously digested with Sal I and Not I) was mixed with 20  $\mu$ g of the purified cDNA, 1X ligase buffer, and 1 ml of T4 DNA  
5 ligase at 16°C for 20 hours.

The product of this ligation was precipitated and introduced into *E. coli* by electroporation, after which transformed bacterial cells were grown in 5 ml SOC medium at 37°C for 1 hour and then frozen at -80°C in 10% glycerol. This constituted the primary Tmrl1 library. Plasmid DNA from the primary Tmrl1  
10 cDNA library was prepared from pools of 50,000 colonies grown on LB/agar plates using standard procedures. Ten pools were prepared and plasmid DNA was isolated from the pools using Qiagen maxi prep kits.

The plasmid DNA that was recovered was subsequently introduced into NIH 3T3 cells by calcium phosphate transfection. In each reaction 100 ng of  
15 plasmid DNA from each pool was used to transfect approximately  $2 \times 10^5$  cells in one 35 mm plate. After 24 hours, the cells were then split into five 100 mm plates, grown in normal medium for one day, and then grown in low serum medium for 13 days. Approximately 4000 total colonies were obtained following growth in the low serum media.

20 Signal peptide-enriched regenerating cDNA molecules were recovered from the transfected cells as follows. Cells were released from the plates by the addition of 2 ml trypsin-EDTA and incubation at 37°C for 5 minutes, followed by gentle swirling. Released cells were transferred to 50 ml conical tubes with 2 ml of fetal calf serum and centrifuged at 1000 rpm for 5 minutes to pellet the cells.  
25 Cell pellets of no more than 1 gram were lysed with 20 ml of TRIZOL reagent (BRL), homogenized for 30 seconds, and then extracted with 4 ml of chloroform. The tubes were centrifuged at 4000 rpm for 30 minutes and the aqueous phase was transferred to a new tube. RNA was precipitated by adding 10 ml isopropanol, mixing, and then centrifuging for 30 minutes at 4200 rpm. The RNA pellet was  
30 washed with 10 ml of 70% ethanol, briefly dried, and then the pellet was

- 78 -

resuspended in 0.5 ml TE buffer. Poly A+ RNA was prepared by using a commercially available mRNA purification kit (Pharmacia). After eluting poly A+ RNA from the column in 750 µl of TE buffer, the sample was then ethanol precipitated in two 1.5 ml microcentrifuge tubes by adding 40 µl of sample buffer and 1 ml of ethanol and incubating overnight at -70°C.

The cDNA inserts of positive clones were rescued by RT-PCR as follows. First strand synthesis was performed using the SuperScript™ preamplification system (BRL). A mixture containing 15 µg of poly A+ RNA from selected 3T3 colonies and 15 µl (2 µM) of vector primer (5'-A-A-T-C-C-G-A-T-G-C-C-C-A-C-G-T-T-G-C-A-G-T-A-3'; SEQ ID NO: 35) was prepared and then incubated at 70°C for 10 minutes followed by equilibration at 50°C. A premixture containing 2.5 µl 10X buffer, 2.5 µl 25 mM MgCl<sub>2</sub>, 1.3 µl 10 mM dNTPs, and 2.5 µl 0.1 M dithiothreitol was then added to the poly A+ RNA/primer mixture, after which 1.2 µl of reverse transcriptase was added and the reaction incubated at 50°C for 1 hour. The reaction was stopped by heating at 70°C for 15 minutes.

Following first strand synthesis, RNA was digested with 1 µl RNase H at 37°C for 20 minutes. PCR was performed using Pfu polymerase (Perkin Elmer) as follows. In a total volume of 100 µl, 2 µl of the first strand reaction was added to 1X Pfu buffer, 0.5 µM each of the amplification primers (5'-A-A-A-A-T-C-T-T-A-G-A-C-C-G-A-C-G-A-C-T-G-T-G-T-T-T-3'; SEQ ID NO: 36; and 5'-G-A-G-T-C-T-C-C-G-C-A-G-C-C-T-T-T-T-G-A-G-G-3'; SEQ ID NO: 37), 0.2 mM dNTPs, 5% DMS, and 2.5 units of Pfu polymerase. The amplification reaction was performed at 95°C for 1 minute for 1 cycle; 95°C for 30 seconds, 66°C for 45 seconds, and 72°C for 30 cycles; and 72°C for 10 minutes for 1 cycle. PCR products were purified by phenol/chloroform extraction followed by ethanol precipitation, and then were digested with Not I and Sal I. Small digestion products and PCR primers were removed from the reaction using SizeSep 400 Spun columns.

Clones identified in the primary screen were subsequently analyzed in a secondary secretion assay. This secondary screening technique utilized a vector



- 79 -

containing a truncated placental alkaline phosphatase (PLAP), in which the native signal peptide had been removed, as a secretion reporter gene. Heterologous cDNA fragments were tested for the presence of signal peptide secretion sequences by inserting individual sequences immediately upstream of the truncated PLAP gene and then transfecting COS7 cells with the test constructs. Inserted cDNA sequences encoding a signal peptide, when inserted in frame with the PLAP reporter sequence, lead to the formation of a fusion protein that can be secreted from the transfected cells.

The PLAP reporter construct was generated as follows. Human placenta RNA was subjected to RT/PCR amplification under standard conditions to generate a DNA fragment encoding a truncated human PLAP protein. The RNA was transcribed by reverse transcriptase using oligo d(T) as primer, and then PCR amplified with PLAP specific primers to produce a DNA product which encodes a PLAP sequence corresponding to amino acids 22 to 536 (Millan, *J. Biol. Chem.* 261(7):3112-15 (1986)). The PLAP amplification primers (5'-A-C-T-G-G-C-G-G-C-C-G-C-A-G-G-C-A-T-C-A-T-C-C-C-A-G-T-T-G-A-G-G-A-G-3'; SEQ ID NO: 38; and 5'-A-C-T-G-G-T-C-A-C-T-C-G-A-G-G-G-T-A-C-C-T-T-A-G-C-T-A-G-C-C-C-C-G-G-G-3'; SEQ ID NO: 39) were designed to contain Not I and Kpn I restriction sites, respectively, in order to facilitate the ligation of the PLAP fragment into the pcDNA3.1 vector (generating the vector pcDNA3.1/PLAP).

The Tmrl1 secondary library was constructed in pcDNA3.1/PLAP as follows. The cDNA inserts from clones in the primary Tmrl1 library were recovered from 3T3 cell colonies using the PCR amplification primers and conditions noted above. Recovered PCR products were then ligated into the pcDNA3.1/PLAP vector which had been digested with the Xho I and Not I restriction enzymes. Ligation products were then transformed into the *E. Coli* strain DH10B to generate the secondary Tmrl1 library.

To assay heterologous cDNA fragments for the presence of signal peptide secretion sequences individual colonies were first selected from a low density plating of the secondary Tmrl1 library on agar plates and placed into the wells of a

- 80 -

standard 96-well plate containing bacterial growth medium. The cultures were then grown to saturation and plasmid DNA was prepared from each culture by standard procedures. Plasmid DNA prepared as such was used to transfect COS7 cells as described below.

5 COS7 cells were seeded in 96-well flat-bottom plates at a concentration of 6000 cells per well in 200  $\mu$ l of a growth medium consisting of DMEM-HG, 10% fetal bovine serum and 1X PSG (penicillin, streptomycin and gentamycin). Following seeding with COS7 cells, the plates were incubated for 24 hours at 37°C in a CO<sub>2</sub> incubator. The cells in each well were subsequently transfected  
10 with 500 ng of plasmid DNA recovered from selected secondary Tmrl1 library clones using Superfect reagent (Qiagen). Transfection reactions were allowed to proceed for two hours, after which the transfected cells were washed with 200  $\mu$ l of phenol red-free, serum-free DMEM and then incubated with 100  $\mu$ l of phenol red-free, serum-free DMEM and glutamate for 24 hours at 37°C in a CO<sub>2</sub>  
15 incubator. Following incubation, 100  $\mu$ l of a solution containing 200  $\mu$ M 4-methylumbelliferyl phosphate (Molecular Probes) in 1M diethanolamine, 10 mM homo-arginine, 1 mM MgCl<sub>2</sub>, and 1 mg/ml BSA was added to each well, and incubation continued for 1 hour at 37°C. The product of the alkaline phosphatase reaction was then read in a fluorometer at 360/460 nm.

20 A single clone (tmrl1-00001-e9), yielding an increased fluorescence readout in the PLAP assay and possessing a computer-predicted signal peptide sequence and homology to the FGF family, was identified in the secondary Tmrl1 library screen. This clone was subsequently used as a probe to isolate a full-length cDNA for murine FGF-like polypeptide from a mouse liver cDNA library.  
25 A mouse liver full-length cDNA library was constructed using standard techniques. Essentially, oligo d(T) was used to prime first strand synthesis from mRNA isolated from regenerating mouse liver and full-length cDNA sequences were then cloned into the pSPORT (Gibco BRL) vector using the Superscript Plasmid System for cDNA Synthesis and Plasmid Cloning (Gibco BRL).

- 81 -

In a primary screen of the mouse liver cDNA library, 50,000 colonies were plated on LB/Ampicillin plates and then transferred to nitrocellulose filters. The filters were screened with a mixture of probes derived from clones identified in the kFGF signal trap and secretion assay screenings. This pool of probes included  
5 a 339 bp Not I-Xba I fragment isolated from the tmrl1-00001-e9 clone. The filters were first prehybridized for 2 hours at 42°C in a hybridization solution consisting of 50% formamide, 5X Denhardt's, 5X SSC, 0.5% SDS, and 100 µg/ml salmon sperm DNA. Following prehybridization, the filters were incubated with the <sup>32</sup>P-dCTP labeled probes overnight at 42°C in fresh hybridization solution. The filters  
10 were then washed with 0.1X SSC/0.1% SDS at 65°C and were subsequently analyzed by autoradiography overnight at -80°C. From this initial screen, 92 positive clones were identified.

The positive clones isolated in the primary screen were pooled and rescreened with the 339 bp tmrl1-00001-e9 fragment alone. In this secondary  
15 screening, 6000 colonies derived from the clone pool were plated on LB/Ampicillin plates and then transferred to nitrocellulose filters. The hybridization conditions used in the secondary screen were the same as those used for the primary screen. Three individual clones (1E, 1E-4, and 1E-6) were identified in the secondary screen.

20 Restriction digestion of the three individual clones identified in the secondary screen indicated that each contained a 1.6 kb insert. Primers corresponding to the 5' (5'-T-G-G-A-A-T-G-G-A-T-G-A-G-A-T-C-T-A-G-A-G-3'; SEQ ID NO: 7) and 3' (5'-C-T-A-G-A-T-T-C-A-G-G-A-A-G-A-G-T-C-A-3'; SEQ ID NO: 8) ends of the coding sequence encoded by this insert were designed  
25 following partial sequence analysis of clone 1E. These primers were used in a polymerase chain reaction (PCR) amplification of clone 1E plasmid DNA. The amplification reaction was performed at 94°C for 1 minute for 1 cycle; 94°C for 15 seconds and 65°C for 1.5 minutes for 35 cycles; and 72°C for 10 minutes for 1 cycle. A PCR product of approximately 650 bp was obtained following  
30 amplification.

- 82 -

Following purification from a 1% agarose gel using a Qiagen gel extraction kit, the PCR product was sequenced. Sequence analysis of this amplification product indicated that the cDNA clones from which the PCR primers were derived contained a gene comprising a 630 bp open reading frame encoding a protein of 210 amino acids. Figure 1 illustrates the nucleotide sequence of the murine FGF-like gene (SEQ ID NO: 3) and the deduced amino acid sequence of murine FGF-like protein (SEQ ID NO: 4). Subsequent sequence analysis of clone 1E also established that the open reading frame of the cDNA clones identified in the secondary screening was interrupted by two intron sequences. Computer analysis using the FASTA program of the Swissprot database indicated that this open reading frame encoded a polypeptide that is most closely related (39% identical) to FGF-6 (Figures 3A-3D).

Computer analysis, using the SIGNALP program (Center for Biological Sequence Analysis, The Technical University of Denmark), also indicated that the murine FGF-like polypeptide possessed a potential signal peptide at its amino terminus (M-E-W-M-R-S-R-V-G-T-L-G-L-W-V-R-L-L-L-A-V-F-L-L-G-V-Y-Q-A; SEQ ID NO: 40; as underlined in Figure 1). The initial translation product of murine FGF-like polypeptide has a calculated molecular weight of 23,237, not including possible post-translational modifications. After removal of the predicted 29 amino acid signal peptide sequence, the remaining predicted mature protein of 181 amino acids has a calculated molecular weight of 19,876. No predicted N-linked glycosylation sites were identified in the protein and the protein does not possess any dibasic protease processing sites.

#### 25 Example 2: Cloning of the Human FGF-like Polypeptide Gene

Generally, materials and methods as described in Sambrook et al. *supra* were used to clone and analyze the gene encoding human FGF-like polypeptide.

Sequences encoding the human FGF-like polypeptide were isolated by screening a human cDNA library with a probe derived from the murine FGF-like polypeptide gene. A 460 bp probe was generated by PCR amplification of murine

- 83 -

FGF-like polypeptide cDNA using the following primers: 5'-T-G-G-A-A-T-G-G-A-T-G-A-G-A-T-C-T-A-G-A-G-3'; SEQ ID NO: 7) and 3' (5'-C-A-T-T-G-C-G-G-C-C-G-C-T-C-A-A-G-A-T-G-C-A-A-A-A-C-G-C-A-G-T-G-3'; SEQ ID NO: 9) in reactions containing <sup>32</sup>P-dCTP. The amplification reaction was performed at 5 94°C for 1 minute for 1 cycle; 94°C for 15 seconds and 65°C for 1.5 minutes for 35 cycles; and 72°C for 10 minutes for 1 cycle.

The 460 bp murine probe was used to isolate a full-length cDNA for human FGF-like polypeptide from a human liver cDNA library. A human liver full-length cDNA library was constructed using standard techniques. Essentially, 10 oligo d(T) was used to prime first strand synthesis from mRNA obtained from Clontech (Palo Alto, CA) and full-length cDNA sequences were then cloned into the pSPORT (Gibco BRL) vector using the Superscript Plasmid System for cDNA Synthesis and Plasmid Cloning (Gibco BRL).

In a primary screen of the human liver cDNA library, 550,000 colonies 15 were plated on LB/Ampicillin plates and then transferred to nitrocellulose filters. The filters were first prehybridized for 30 minutes at 60°C in ExpressHyb solution (Clontech), and then were incubated with the <sup>32</sup>P-dCTP labeled murine FGF-like cDNA probe overnight at 60°C in fresh ExpressHyb solution. Following hybridization, the filters were washed twice for 30 minutes at room temperature in 20 2X SSC/0.1% SDS, twice for 30 minutes at 60°C in 1X SSC/0.1% SDS, twice for 30 minutes at 65°C in 0.1XSSC/0.1% SDS, and then the filters were analyzed by autoradiography overnight at -80°C. Positive clones identified in the primary screen were then rescreened, and four independent clones were recovered following the secondary screen. Plasmid DNA for these four clones was prepared 25 and sequenced.

Sequence analysis indicated that the four clones contained inserts of either 1.2 kb or 1.8 kb. The 1.2 kb cDNA insert contained a gene comprising a 627 bp open reading frame encoding a protein of 209 amino acids. Figures 2A-2B illustrate the nucleotide sequence of the human FGF-like gene (SEQ ID NO: 1) 30 and the deduced amino acid sequence of human FGF-like protein (SEQ ID NO:



- 84 -

2). While the 1.8 kb cDNA insert contained the same open reading frame as encoded by the 1.2 kb insert, the open reading frame of this insert was additionally interrupted by an intron corresponding in location to the second intron found in some of the murine cDNA clones isolated in Example 1.

5        Figures 3A-3D illustrate the amino acid sequence alignment of human FGF-like protein, murine FGF-like protein, and other members of the FGF family. The human FGF-like polypeptide is 76% identical to the murine FGF-like protein. Computer analysis, using the SIGNALP program (Center for Biological Sequence Analysis, The Technical University of Denmark), indicated that the human FGF-  
10    like polypeptide also possessed a potential signal peptide at its amino terminus (M-D-S-D-E-T-G-F-E-H-S-G-L-W-V-S-V-L-A-G-L-L-L-G-A-C-Q-A; SEQ ID NO: 41; as underlined in Figures 2A-2B). The initial translation product of human FGF-like polypeptide has a calculated molecular weight of 22,284, not including possible post-translational modifications. After removal of the  
15    predicted 28 amino acid signal peptide sequence, the remaining predicted mature protein of 181 amino acids has a calculated molecular weight of 19,395. No predicted N-linked glycosylation sites were identified in the protein and the protein does not possess any dibasic protease processing sites.

### 20    Example 3: FGF-like mRNA Expression

Expression of murine FGF-like mRNA was examined on a murine multiple tissue Northern blot (Clontech) using a <sup>32</sup>P-dCTP labeled murine FGF-like cDNA probe. The probe consisted of a 391 bp fragment isolated from the tmrl1-00001-e9 clone by restriction digestion with Xba I and Not I. The blot was  
25    first prehybridized for 2 hours at 42°C in a hybridization solution consisting of 50% formamide, 5X Denhardt's, 6X SSC, 0.5% SDS, and 100 µg/ml salmon sperm DNA. Following prehybridization, the filters were incubated with the <sup>32</sup>P-dCTP labeled murine FGF-like cDNA probe overnight at 42°C in fresh hybridization solution. The filters were then washed with 0.1X SSC/0.1% SDS at  
30    65°C and were subsequently analyzed by autoradiography overnight at -80°C.

- 85 -

Two transcripts, of approximately 1.35 kb and 1.8 kb, were detected in murine liver (Figure 4A). The 1.35 kb transcript showed the predominant expression.

Expression of human FGF-like mRNA was examined on a Human RNA Master Blot™ (Clontech) and a human multiple tissue Northern blot (Clontech) using a <sup>32</sup>P-dCTP labeled human FGF-like cDNA probe. The probe consisted of a 660 bp PCR product derived from the human FGF-like protein coding region using the primers: 5'-C-T-A-C-TA-A-A-G-C-T-T-C-C-A-C-C-A-T-G-G-A-C-T-C-G-G-A-C-G-A-G-A-C-C-G-3'; SEQ ID NO: 12; and 5'-A-T-T-C-A-T-G-C-G-G-C-C-G-C-G-G-A-A-G-C-G-T-A-G-C-T-G-G-G-G-C-T-T-C-3'; SEQ ID NO: 13, and the human cDNA clone described above as a template. The amplification reaction was performed at 94°C for 1 minute for 1 cycle; 94°C for 15 seconds, 60°C for 15 seconds, and 72°C for 1 minute for 35 cycles; and 72°C for 10 minutes for 1 cycle. The blots were first prehybridized for 1 hour at 65°C in ExpressHyb solution (Clontech), and then were incubated with the <sup>32</sup>P-dCTP labeled human FGF-like cDNA probe overnight at 65°C in fresh ExpressHyb solution. Following hybridization, the filters were washed twice for 30 minutes at room temperature in 2X SSC/0.1% SDS, twice for 30 minutes at 65°C in 0.1X SSC/ 0.1% SDS, and then the filters were analyzed by autoradiography overnight at -80°C. Two transcripts, of approximately 1.2 kb and 2 kb, were detected in human liver on the multiple tissue Northern blot 9 (Figure 4B). Strong expression in adult liver and weak expression in lung and fetal liver was detected on the Human RNA Master Blot™ (Figure 4C).

#### Example 4: FGF-like mRNA *in situ* analysis

*In situ* hybridization was performed with a 648 bp antisense RNA probe spanning the coding region of murine FGF-like polypeptide. The probe was transcribed from a linearized pCR2.1 TOPO plasmid containing the FGF-like cDNA insert using T7 RNA polymerase and <sup>33</sup>P-UTP.

A panel of normal embryonic and adult mouse tissues were fixed in 4% paraformaldehyde, embedded in paraffin, and sectioned at 5 μm. Prior to *in situ*

- 86 -

hybridization, tissues were permeabilized with 0.2M HCl, followed by digestion with Proteinase K, and acetylation with triethanolamine and acetic anhydride. Sections were hybridized with the <sup>33</sup>P-labeled riboprobe overnight at 55°C, then subjected to a high stringency wash in 0.1 X SSC at 60°C. Slides were dipped in  
5 Kodak NTB2 emulsion, exposed at 4°C for 2-3 weeks, developed, and counterstained with hematoxylin and eosin. Sections were examined with darkfield and standard illumination to allow simultaneous evaluation of tissue morphology and hybridization signal.

The strongest overall expression was noted in the pancreas with a strong  
10 signal being detected over the islets and a lower, more diffuse signal over the acinar portion of the pancreas. The liver displayed a moderate level of diffuse signal, indicative of moderate hepatocellular expression of FGF-like polypeptide. Significant signal was also present over spermatogonia within the seminiferous tubules in the testis and over cells in the thymic medulla. A low level of diffuse  
15 signal was detected in kidney, spleen, pituitary, and white and brown adipose tissue.

#### Example 5: Production of FGF-like Polypeptides in Mammalian Cells

Both the human and murine FGF-like polypeptides were expressed as  
20 fusion proteins with a human immunoglobulin IgG heavy chain Fc region at their carboxyl terminus. Template DNA sequences encoding human or murine FGF-like polypeptide were amplified by PCR using primers corresponding to the 5' and 3' ends of the sequence (Table II). The resulting PCR products corresponded to the coding region of either the human or murine FGF-like polypeptide, minus the  
25 translation termination codon. In addition, the primers were designed to incorporate a Hind III restriction endonuclease site at the 5' end of the PCR product and a Not I site at the 3' end of the product.

Table II

30 Primers Used in Recombinant Protein Expression

- 87 -

Murine 5' 5'-CTACTAAAGCTTCCACCATGGAATGGATGAGATCTAG-3' (SEQ ID NO: 10)

Murine 3' 5'-ATTCATGCGGCCGCGGACGCATAGCTGGGGCTT-3' (SEQ ID NO: 11)

Human 5' 5'-CTACTAAAGCTTCCACCATGGACTCGGACGAGACCG-3' (SEQ ID NO: 12)

Human 3' 5'-ATTCATGCGGCCGCGGAAGCGTAGCTGGGGCTTC-3' (SEQ ID NO: 13)

5

The human and murine FGF-like protein-Fc fusion constructs were generated by first cleaving the PCR products with Hind III and Not I and then ligating the fragments in frame to a DNA fragment encoding the HFc chain of human immunoglobulin IgG. The FGF-like protein-Fc insert was then ligated into the pCEP4 mammalian expression vector (Invitrogen). These ligations were transformed into the *E. coli* strain DH10 by electroporation and transformants selected for ampicillin resistance. Following sequence analysis of selected transformants, large-scale plasmid stocks were prepared for tissue culture transfection. Plasmid DNA for selected ampicillin resistant colonies was prepared and sequenced to confirm that the clone contained the desired insert.

To conduct functional studies on FGF-like protein, human or murine FGF-like protein-Fc fusion expression constructs were introduced into 293-EBNA (Invitrogen) cells using SuperFect™ transfection reagent (Qiagen). The conditioned medium was harvested 48 hours after transient transfection. Western blot analysis, using an anti-human Fc antibody, confirmed that the conditioned media contained human or murine FGF-like/Fc fusion polypeptides.

Conditioned medium was purified by affinity chromatography as described below. The medium was first passed through a 0.2 µm filter. Protein A columns (Pharmacia) were equilibrated with ImmunoPure Gentle Binding Buffer (Pierce, Rockford, IL), and then loaded with the filtered medium. The column was washed with ImmunoPure Gentle Binding Buffer until the absorbance at 280 nm reached a baseline. FGF-like/Fc protein was eluted from the column with ImmunoPure Gentle Binding Buffer. Fractions containing FGF-like/Fc protein were pooled, dialyzed in Tris-buffered saline (TBS) followed by Phosphate-buffered saline (PBS), and stored at 4°C. Gel electrophoretic analysis confirmed

- 88 -

that the pooled fractions contained a purified protein of the expected molecular weight of about 60 kD.

Example 6: Production and Analysis of Transgenic Mice Overexpressing FGF-like Polypeptide

Transgenic mice overexpressing mouse FGF-like polypeptide (SEQ ID NO: 4) from the human apolipoprotein E promoter were generated as previously described (see Simonet et al., 1997, Cell 89:309-19). Seven mice (four males and three females) which were transgenic for the murine FGF-like gene (SEQ ID NO: 3) and five non-transgenic littermates (two males and three females) underwent necropsy and pathological analysis at 6-8 weeks of age. All of the mice were injected with 50 mg/kg of BrdU one hour prior to harvest, then were radiographed and sacrificed. Body and selected organ weights were measured, blood was drawn for hematology and serum chemistries, and organs were harvested for histological analysis and BrdU labeling.

All of the transgenic mice were under 20 gm body weight while all of the non-transgenic mice were over 20 gm body weight ( $p < 0.0001$ ). In addition, transgenic mice had statistically significantly lower liver ( $p = 0.0011$ ) and spleen weights ( $p = 0.0039$ ) and a higher thymic ( $p = 0.0118$ ) weight (as a percent of body weight) than their non-transgenic counterparts. For the transgenic mice, body weight was 67% of wild type. Liver, spleen, and thymus weights, as a percentage of body weight, were, respectively, 85%, 63% and 170% of wild type. All female transgenic mice also had small uteri and oviducts and ovaries that lacked corpora lutea and exhibited little follicular development. In summary, transgenic mice had a phenotype that is best characterized as stunted growth with small livers, spleens, and poorly developed ovaries, and enlarged (probably not involuted) thymuses. These changes are most consistent with overexpression of FGF-like polypeptide in the transgenics causing inhibited or delayed maturation in comparison with their age-matched non-transgenic littermates.



Example 7: Analysis of One-Year-Old FGF-like Polypeptide Transgenic Mice

Necropsy and pathological analysis was performed on one-year-old FGF-like polypeptide transgenics (five males and five females) and non-transgenic (three males and two females) littermates that had been produced in the experiments described in Example 6. FGF-like polypeptide transgenics continued to demonstrate an abnormal phenotype generally characterized as inhibited or delayed maturation. These observed effects included reduced body weight (48% of wild type) and, in the females, poorly developed ovaries with lack of significant follicular development. Liver, spleen, and thymus weights as a percentage of body weight had normalized to that found in the non-transgenic littermates. Several of the one-year-old non-transgenic control mice were found to be obese, and at least one of the controls exhibited changes consistent with the development of type II diabetes. However, none of the one-year-old transgenic mice were obese or showed any evidence of developing diabetes. Thus, it appears that FGF-like polypeptide transgenics do not develop at least some of the age related changes commonly seen in mice as they age and, indeed, that the FGF-like gene of this invention may help retard the aging process.

These findings are significant and support the conclusion that the FGF-like polynucleotides and polypeptides of the present invention may be useful for the treatment or diagnosis of age-related diseases, disorders, or conditions. By way of illustration, such diseases, disorders or conditions may include, without limitation, atherosclerosis, hypercholesterolemia, diabetes, obesity, stroke, osteoporosis, osteoarthritis, degenerative joint disease, muscle atrophy, sarcopenia, decreased lean body mass, baldness, wrinkles, increased fatigue, decreased stamina, decreased cardiac function, immune system dysfunction, cancer, Parkinson's disease, senile dementia, Alzheimer's disease, and decreased cognitive function. More generally, the molecules of the present invention may be applicable for enhancing or increasing life-span.

- 90 -

While the present invention has been described in terms of the preferred embodiments, it is understood that variations and modifications will occur to those skilled in the art. Therefore, it is intended that the appended claims cover all such equivalent variations that come within the scope of the invention as  
5 claimed.

- 91 -

## WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:
  - 5 (a) the nucleotide sequence as set forth in SEQ ID NO: 1 or SEQ ID NO: 3;
  - (b) the nucleotide sequence of the DNA insert in ATCC Deposit No. PTA-626;
  - (c) a nucleotide sequence encoding the polypeptide as set forth in SEQ  
10 ID NO: 2 or SEQ ID NO: 4;
  - (d) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a)-(c); and
  - (e) a nucleotide sequence complementary to any of (a)-(c).
- 15 2. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:
  - (a) a nucleotide sequence encoding a polypeptide that is at least about 80 percent identical to the polypeptide as set forth in SEQ ID NO: 2 or SEQ ID NO: 4, wherein the encoded polypeptide activates one or more FGF receptors,  
20 regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;
  - (b) a nucleotide sequence encoding an allelic variant or splice variant of the nucleotide sequence as set forth in SEQ ID NO: 1, SEQ ID NO: 3, or (a);
  - 25 (c) a region of the nucleotide sequence of SEQ ID NO: 1, SEQ ID NO: 3, (a), or (b) encoding a polypeptide fragment of at least about 25 amino acid residues, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a

- 92 -

role in liver or pancreas chemotaxis, has an oncogenic activity, or serves as an antigen for generating antibodies;

(d) a region of the nucleotide sequence of SEQ ID NO: 1, SEQ ID NO: 3, or any of (a) - (c) comprising a fragment of at least about 16 nucleotides;

5 (e) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (d); and

(f) a nucleotide sequence complementary to any of (a) - (d).

3. An isolated nucleic acid molecule comprising a nucleotide  
10 sequence selected from the group consisting of:

(a) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one conservative amino acid substitution, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types  
15 following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;

(b) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one amino acid insertion, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of  
20 cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;

(c) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one amino acid deletion, wherein the encoded polypeptide  
25 activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;

(d) a nucleotide sequence encoding a polypeptide as set forth in SEQ  
30 ID NO: 2 which has a carboxyl- and/or amino-terminal truncation, wherein the

- 93 -

encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;

5 (e) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, carboxyl-terminal truncation, and amino-terminal truncation, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and  
10 differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;

(f) a region of the nucleotide sequence of any of (a) - (e) comprising a fragment of at least about 16 nucleotides;

15 (g) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (f); and

(h) a nucleotide sequence complementary to any of (a) - (e).

20 4. A vector comprising the nucleic acid molecule of Claims 1, 2, or 3.

5. A host cell comprising the vector of Claim 4.

6. The host cell of Claim 5 that is a eukaryotic cell.

25 7. The host cell of Claim 5 that is a prokaryotic cell.

8. A process of producing a FGF-like polypeptide comprising culturing the host cell of Claim 5 under suitable conditions to express the polypeptide, and optionally isolating the polypeptide from the culture.

30



- 94 -

9. A polypeptide produced by the process of Claim 8.

10. The process of Claim 8, wherein the nucleic acid molecule comprises promoter DNA other than the promoter DNA for the native FGF-like polypeptide operatively linked to the DNA encoding the FGF-like polypeptide.

11. The isolated nucleic acid molecule according to Claim 2 wherein the percent identity is determined using a computer program selected from the group consisting of GAP, BLASTN, FASTA, BLASTA, BLASTX, BestFit, and the Smith-Waterman algorithm.

12. A process for determining whether a compound inhibits FGF-like polypeptide activity or FGF-like polypeptide production comprising exposing a cell according to Claims 5, 6, or 7 to the compound and measuring FGF-like polypeptide activity or FGF-like polypeptide production in said cell.

13. An isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence as set forth in SEQ ID NO: 2 or SEQ ID NO: 4; and
- (b) the amino acid sequence encoded by the DNA insert of ATCC Deposit No. PTA-626.

14. An isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence as set forth in either SEQ ID NO: 5 or SEQ ID NO: 6, optionally further comprising an amino-terminal methionine;
- (b) an amino acid sequence for an ortholog of either SEQ ID NO: 2 or SEQ ID NO: 4, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or

- 95 -

pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;

(c) an amino acid sequence that is at least about 80 percent identical to the amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 4, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;

(d) a fragment of the amino acid sequence set forth in either SEQ ID NO: 2 or SEQ ID NO: 4 comprising at least about 25 amino acid residues, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, has an oncogenic activity, or serves as an antigen for generating antibodies; and

(e) an amino acid sequence for an allelic variant or splice variant of either the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 4; the amino acid sequence encoded by the DNA insert of ATCC Deposit No. PTA-626; (a), (b), or (c).

20

15. An isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

(a) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 4 with at least one conservative amino acid substitution, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;

(b) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 4 with at least one amino acid insertion, wherein the encoded

30

- 96 -

polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity;

5 (c) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 4 with at least one amino acid deletion, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas  
10 chemotaxis, or has an oncogenic activity;

(d) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 4 which has a C- and/or N- terminal truncation, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types  
15 following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity; and

(e) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 4 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-  
20 terminal truncation, and N-terminal truncation, wherein the encoded polypeptide activates one or more FGF receptors, regulates the growth and differentiation of cells within the liver or pancreas, regulates other cell types following secretion from the liver or pancreas, plays a role in liver or pancreas chemotaxis, or has an oncogenic activity.

25

16. An isolated polypeptide encoded by the nucleic acid molecule of any of Claims 1, 2, or 3.

17. The isolated polypeptide according to Claim 14 wherein the  
30 percent identity is determined using a computer program selected from the group

- 97 -

consisting of GAP, BLASTP, FASTA, BLASTA, BLASTX, BestFit, and the Smith-Waterman algorithm.

18. An antibody produced by immunizing an animal with a peptide  
5 comprising an amino acid sequence of SEQ ID NO: 2 or SEQ ID NO: 4.

19. An antibody or fragment thereof that specifically binds the polypeptide of any of Claims 13, 14, or 15.

10 20. The antibody of Claim 19 that is a monoclonal antibody.

21. A hybridoma that produces a monoclonal antibody that binds to a peptide comprising an amino acid sequence of SEQ ID NO: 2 or SEQ ID NO: 4.

15 22. A method of detecting or quantitating the amount of FGF-like polypeptide using the anti-FGF-like antibody of any of Claims 18, 19, or 20 or a fragment thereof.

23. A composition comprising the polypeptide of any of Claims 13, 14,  
20 or 15 and a pharmaceutically acceptable formulation agent.

24. The composition of Claim 23 wherein the pharmaceutically acceptable formulation agent is a carrier, adjuvant, solubilizer, stabilizer, or anti-oxidant.

25 25. The composition of Claim 22 wherein the polypeptide comprises the mature amino acid sequence as set forth in SEQ ID NO: 2.

26. A polypeptide comprising a derivative of the polypeptide of any of  
30 Claims 13, 14, or 15.

- 98 -

27. The polypeptide of Claim 26 which is covalently modified with a water-soluble polymer.

5           28. The polypeptide of Claim 27 wherein the water-soluble polymer is selected from the group consisting of polyethylene glycol or dextran.

29. A fusion polypeptide comprising the polypeptide of any of Claims 13, 14, or 15 fused to a heterologous amino acid sequence.

10

30. The fusion polypeptide of Claim 29 wherein the heterologous amino acid sequence is an IgG constant domain or fragment thereof.

31. A method for treating, preventing or ameliorating a medical  
15 condition comprising administering to a patient the polypeptide of any of Claims 13, 14, or 15 or the polypeptide encoded by the nucleic acid of any of Claims 1, 2, or 3.

32. A method for treating, preventing or ameliorating a medical  
20 condition comprising administering to a patient an agonist or antagonist of the biological activity of the polypeptide of any of Claims 13, 14, or 15 or of the polypeptide encoded by the nucleic acid of any of Claims 1, 2, or 3.

33. The method of either Claims 31 or 32 wherein the medical  
25 condition being treated, prevented, or ameliorated is cirrhosis or other toxic insult of the liver; inflammatory bowel disease, mucositis, Crohn's disease, or other gastrointestinal abnormality; diabetes; obesity; neurodegenerative diseases; wounds; damage to the corneal epithelium, lens, or retinal tissue; damage to renal tubules as a result of acute tubular necrosis; hematopoietic cell reconstitution  
30 following chemotherapy; wasting syndromes (for example, cancer associated



- 99 -

cachexia), multiple sclerosis, myopathies; short stature, delayed maturation, excessive growth (for example, acromegaly), premature maturation; alopecia; diseases or abnormalities of androgen target organs; infantile respiratory distress syndrome, bronchopulmonary dysplasia, acute respiratory distress syndrome, or  
5 other lung abnormalities; tumors of the eye or other tissues; atherosclerosis; hypercholesterolemia; diabetes; obesity; stroke; osteoporosis; osteoarthritis; degenerative joint disease; muscle atrophy; sarcopenia; decreased lean body mass; baldness; wrinkles; increased fatigue; decreased stamina; decreased cardiac function; immune system dysfunction; cancer; Parkinson's disease; senile  
10 dementia; Alzheimer's disease; and decreased cognitive function.

34. A method for modulating body growth or maturation or life-span comprising administering to a patient the polypeptide of any of Claims 13, 14, or 15 or the polypeptide encoded by the nucleic acid of any of Claims 1, 2, or 3 or an  
15 agonist or antagonists of the activity thereof.

35. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:

(a) determining the presence or amount of expression of the  
20 polypeptide of any of Claims 13, 14, or 15 or the polypeptide encoded by the nucleic acid molecule of any of Claims 1, 2, or 3 in a sample; and

(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or amount of expression of the polypeptide.

25

36. A device, comprising:

(a) a membrane suitable for implantation; and

(b) cells encapsulated within said membrane, wherein said cells secrete a protein of any of Claims 13, 14, or 15;

30 said membrane being permeable to said protein product and impermeable

- 100 -

to materials detrimental to said cells.

37. A method of identifying a compound which binds to a polypeptide comprising:

5 (a) contacting the polypeptide of any of Claims 13, 14, or 15 with a compound; and

(b) determining the extent of binding of the polypeptide to the compound.

10 38. The method of Claim 37, further comprising determining the activity of the polypeptide when bound to the compound.

39. A method of modulating levels of a polypeptide in an animal comprising administering to the animal the nucleic acid molecule of any of  
15 Claims 1, 2, 3, or 4.

40. A transgenic non-human mammal comprising the nucleic acid molecule of any of Claims 1, 2, 3, or 4.

20 41. A process for determining whether a compound inhibits FGF-like polypeptide activity or FGF-like polypeptide production comprising exposing a transgenic mammal according to Claim 40 to the compound, and measuring FGF-like polypeptide activity or FGF-like polypeptide production in said mammal.

1/10

Fig. 1

atg gaa tgg atg aga tct aga gtt ggg acc ctg gga ctg tgg gtc cga	48
Met Glu Trp Met Arg Ser Arg Val Gly Thr Leu Gly Leu Trp Val Arg	
1 5 10 15	
ctg ctg ctg gct gtc ttc ctg ctg ggg gtc tac caa gca tac ccc atc	96
Leu Leu Leu Ala Val Phe Leu Leu Gly Val Tyr Gln Ala Tyr Pro Ile	
20 25 30	
cct gac tcc agc ccc ctc ctc cag ttt ggg ggt caa gtc cgg cag agg	144
Pro Asp Ser Ser Pro Leu Leu Gln Phe Gly Gly Gln Val Arg Gln Arg	
35 40 45	
tac ctc tac aca gat gac gac caa gac act gaa gcc cac ctg gag atc	192
Tyr Leu Tyr Thr Asp Asp Asp Gln Asp Thr Glu Ala His Leu Glu Ile	
50 55 60	
agg gag gat gga aca gtg gta ggc gca gca cac cgc agt cca gaa agt	240
Arg Glu Asp Gly Thr Val Val Gly Ala Ala His Arg Ser Pro Glu Ser	
65 70 75 80	
ctc ctg gag ctc aaa gcc ttg aag cca ggg gtc att caa atc ctg ggt	288
Leu Leu Glu Leu Lys Ala Leu Lys Pro Gly Val Ile Gln Ile Leu Gly	
85 90 95	
gtc aaa gcc tct agg ttt ctt tgc caa cag cca gat gga gct ctc tat	336
Val Lys Ala Ser Arg Phe Leu Cys Gln Gln Pro Asp Gly Ala Leu Tyr	
100 105 110	
gga tcg cct cac ttt gat cct gag gcc tgc agc ttc aga gaa ctg ctg	384
Gly Ser Pro His Phe Asp Pro Glu Ala Cys Ser Phe Arg Glu Leu Leu	
115 120 125	
ctg gag gac ggt tac aat gtg tac cag tct gaa gcc cat ggc ctg ccc	432
Leu Glu Asp Gly Tyr Asn Val Tyr Gln Ser Glu Ala His Gly Leu Pro	
130 135 140	
ctg cgt ctg cct cag aag gac tcc cca aac cag gat gca aca tcc tgg	480
Leu Arg Leu Pro Gln Lys Asp Ser Pro Asn Gln Asp Ala Thr Ser Trp	
145 150 155 160	
gga cct gtg cgc ttc ctg ccc atg cca ggc ctg ctc cac gag ccc caa	528
Gly Pro Val Arg Phe Leu Pro Met Pro Gly Leu Leu His Glu Pro Gln	
165 170 175	
gac caa gca gga ttc ctg ccc cca gag ccc cca gat gtg ggc tcc tct	576
Asp Gln Ala Gly Phe Leu Pro Pro Glu Pro Pro Asp Val Gly Ser Ser	
180 185 190	
gac ccc ctg agc atg gta gag cct tta cag ggc cga agc ccc agc tat	624
Asp Pro Leu Ser Met Val Glu Pro Leu Gln Gly Arg Ser Pro Ser Tyr	
195 200 205	

**Fig. 2A**

gag	gat	ccag	ccg	aaag	agg	agc	cagg	ccac	tcagg	ccacc	tgagt	ctact	cac	ctg	ggaca	60
act	gga	atct	ggc	acca	aatt	ctaa	acc	act	cag	ctt	ctcc	gag	ctc	acac	ccc	ggagatc 120
ac	ctg	agg	gac	ccg	agc	catt	g	atg	gac	tcg	gac	gag	acc	ggg	ttc	gag cac 171
								Met	Asp	Ser	Asp	Glu	Thr	Gly	Phe	Glu His
								1				5				10
tca	gga	ctg	tgg	gtt	tct	gtg	ctg	gct	ggc	ctt	ctg	ctg	gga	gcc	tgc	219
Ser	Gly	Leu	Trp	Val	Ser	Val	Leu	Ala	Gly	Leu	Leu	Leu	Gly	Ala	Cys	
				15					20					25		
cag	gca	cac	ccc	atc	cct	gac	tcc	agt	cct	ctc	ctg	caa	ttc	ggg	ggc	267
Gln	Ala	His	Pro	Ile	Pro	Asp	Ser	Ser	Pro	Leu	Leu	Gln	Phe	Gly	Gly	
			30					35					40			
caa	gtc	cgg	cag	cgg	tac	ctc	tac	aca	gat	gat	gcc	cag	cag	aca	gaa	315
Gln	Val	Arg	Gln	Arg	Tyr	Leu	Tyr	Thr	Asp	Asp	Ala	Gln	Gln	Thr	Glu	
		45					50					55				
gcc	cac	ctg	gag	atc	agg	gag	gat	ggg	acg	gtg	ggg	ggc	gct	gct	gac	363
Ala	His	Leu	Glu	Ile	Arg	Glu	Asp	Gly	Thr	Val	Gly	Gly	Ala	Ala	Asp	
	60					65					70					
cag	agc	ccc	gaa	agt	ctc	ctg	cag	ctg	aaa	gcc	ttg	aag	ccg	gga	gtt	411
Gln	Ser	Pro	Glu	Ser	Leu	Leu	Gln	Leu	Lys	Ala	Leu	Lys	Pro	Gly	Val	
	75				80					85					90	
att	caa	atc	ttg	gga	gtc	aag	aca	tcc	agg	ttc	ctg	tgc	cag	cgg	cca	459
Ile	Gln	Ile	Leu	Gly	Val	Lys	Thr	Ser	Arg	Phe	Leu	Cys	Gln	Arg	Pro	
			95					100						105		
gat	ggg	gcc	ctg	tat	gga	tcg	ctc	cac	ttt	gac	cct	gag	gcc	tgc	agc	507
Asp	Gly	Ala	Leu	Tyr	Gly	Ser	Leu	His	Phe	Asp	Pro	Glu	Ala	Cys	Ser	
			110					115					120			
ttc	cgg	gag	ctg	ctt	ctt	gag	gac	gga	tac	aat	gtt	tac	cag	tcc	gaa	555
Phe	Arg	Glu	Leu	Leu	Leu	Glu	Asp	Gly	Tyr	Asn	Val	Tyr	Gln	Ser	Glu	
		125					130					135				
gcc	cac	ggc	ctc	ccg	ctg	cac	ctg	cca	ggg	aac	aag	tcc	cca	cac	cgg	603
Ala	His	Gly	Leu	Pro	Leu	His	Leu	Pro	Gly	Asn	Lys	Ser	Pro	His	Arg	
	140					145					150					
gac	cct	gca	ccc	cga	gga	cca	gct	cgc	ttc	ctg	cca	cta	cca	ggc	ctg	651
Asp	Pro	Ala	Pro	Arg	Gly	Pro	Ala	Arg	Phe	Leu	Pro	Leu	Pro	Gly	Leu	
	155				160					165					170	
ccc	ccc	gca	ccc	ccg	gag	cca	ccc	gga	atc	ctg	gcc	ccc	cag	ccc	ccc	699
Pro	Pro	Ala	Pro	Pro	Glu	Pro	Pro	Gly	Ile	Leu	Ala	Pro	Gln	Pro	Pro	
				175					180					185		

3/10

Fig. 2B

```
gat gtg ggc tcc tcg gac cct ctg agc atg gtg gga cct tcc cag ggc 747
Asp Val Gly Ser Ser Asp Pro Leu Ser Met Val Gly Pro Ser Gln Gly
      190              195              200

cga agc ccc agc tac gct tcc tga agccagaggc tgtttactat gacatctcct 801
Arg Ser Pro Ser Tyr Ala Ser
      205              210

ctttatttat taggttat tttttttatt tttcttactt gagataataa 861
agagttccag aggaggataa gaatgagcat gtgtgagtgt ctgagggaag acatggcagc 921
tgttttgtct cccttggccc ggacaatccc ctctacacct cccctcacgt ggtccgaggg 981
tcctggcttc ccactgggccc tcactttttt cttttctttt cttttctttt ttttgagacg 1041
gagtctcgct ctgcactcca gccagggcca cagagcgaga ttccatctca aaaaaataaa 1101
taaataaata aataaataaa tataaaaata aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1161
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1190
```

4/10

Fig. 3A

	1				50
hAgp-26257	-----	-----	-----	-----	-----M
mAgp-26257	-----	-----	-----	-----	-----M
Hfgf14	-----	-----	--MAAAIASG	LIRQKRQARE	QHWDRPSASR
Mfgf14	-----	-----	--MAAAIASG	LIRQKRQARE	QHWDRPSASR
Hfgf12	-----	-----	--MAAAIASS	LIRQKRQARE	SNSDRVSASK
Mfgf13	-----	-----	--MTAAIASS	LIRQKRQARE	R..EKSNAACK
Hfgf5	-----MS	LSFLLLLFFS	HLILSAWAHG	EKRLAPKGQP	GPAATDRNPI
Mfgf5	-----MS	LSLLFLIFCS	HLIHSAWAHG	EKRLTPEGQP	APPRNP GDSS
Hfgf6	MALGQKLFIT	MSRGAGRLQG	TLWALVFLGI	LVGMVVPSPA	GTRA..NNTL
Mfgf6	MALGQRLFIT	MSRGAGRVQG	TLQALVFLGV	LVGMVVPSPA	GARA..NGTL
Hfgf4	-----	-MSGPGTAAV	ALLPAVLLAL	LAPWAGRGA	AAPTAPNGTL
Mfgf4	-----	-MAKRGPTTG	TLLPRVLLAL	VVALADRGTA	...APNGTR
Hfgf3	-----	-----	-----	-----	-----
Mfgf3	-----	-----	-----	-----	-----
Hfgf7	-----	-----	-----	MHKWILTWIL	PTLLYRSCFH
Mfgf7	-----	-----	-----	MRKWILTRIL	PTLLYRSCFH
Hfgf9	-----	-----	-----	MAPLGEVGNV	FGVQDAVPFG
Mfgf9	-----	-----	-----	MAPLGEVGSY	FGVQDAVPFG
Hfgf1	-----	-----	-----	-----	-----
Mfgf1	-----	-----	-----	-----	-----
Hfgf2	-----	-----	-----	-----	-----
Mfgf2	-----	-----	-----	-----	-----
cons	-----	-----	-----	-----	-----

	51				100
hAgp-26257	DSDETGF EHS	GLWVS.VLAG	LLLGACQ AHP	IPDSSPLLQF	GGQVRQRYLY
mAgp-26257	EWMRSRVGTL	GLWVRLLLAV	FLLGVYQAYP	IPDSSPLLQF	GGQVRQRYLY
Hfgf14	RRSSPSKN.R	GLCNGNLVDI	FSKVRIFGLK	KRRLRRQ.DP	QLKGIVTRLY
Mfgf14	RRSSPSKN.R	GLFNGNLVDI	FSKVRIFGLK	KRRLRRQ.DP	QLKGIVTRLY
Hfgf12	RRSSPSKDGR	SLCERHVLGV	FSKVRFCSGR	KRPVRRRPEP	QLKGIVTRLF
Mfgf13	CVSSPSK.GK	TSCDKNKLNV	FSRVKLF GSK	KRR.RRRPEP	QLKGIVTKLY
Hfgf5	GSSSRQSSSS	AMSSSSASSS	PAASLGSGGS	GLEQSSFQWS	PSGRRTGSLY
Mfgf5	GSRGR..SSA	TFSSSSASSP	VAASPGSGGS	GSEHSSFQWS	PSGRRTGSLY
Hfgf6	LD..SRGWGT	LLSRSRAGL.	.AGEI..AGV	NWESG...YL	VGIKRQRRLY
Mfgf6	LD..SRGWGT	LLSRSRAGL.	.AGEI..SGV	NWESG...YL	VGIKRQRRLY
Hfgf4	EAELERRWES	LVALSLARLP	VAAQP..KEA	AVQSGAGDYL	LGIKRLRRLY
Mfgf4	HAELGHGWDG	LVARSLARLP	VAAQP..PQA	AVRSGAGDYL	LGLKRLRRLY
Hfgf3	-MGLIWLLLL	SLLEPGWPAA	GPGARLRRDA	GGRGGVYEHL	GGAPRRRKLY
Mfgf3	-MGLIWLLLL	SLLEPSWPTT	GPGTRLRRDA	GGRGGVYEHL	GGAPRRRKLY
Hfgf7	IICLVGTISL	ACNDMTPEQM	ATNVNCSSPE	RHTRSYDYME	GGDIRVRRLF
Mfgf7	LVCLVGTISL	ACNDMSPEQT	ATSVNCSSPE	RHTRSYDYME	GGDIRVRRLF
Hfgf9	NVPVLPVDSP	VLLSDHLGQS	EAGGLPRGPA	.VTDLDHLK.	.GILRRRQLY
Mfgf9	NVPVLPVDSP	VLLNDHLGQS	EAGGLPRGPA	.VTDLDHLK.	.GILRRRQLY
Hfgf1	-----	-----MAEG	EITTF TALTE	....KFNLPP	GNYYKPKLLY
Mfgf1	-----	-----MAEG	EITTF AALTE	....RFNLPL	GNYYKPKLLY
Hfgf2	-----	-----MAAG	SITTLPALPE	.DGGSGAFPP	GHFKDPKRLY
Mfgf2	-----	-----MAAS	GITSLPALPE	.DGGGA.AFPP	GHFKDPKRLY



5/10

Fig. 3B

	101		150
hAgp-26257	TDDAQQTEAH	LEIREDGTVG	GAADQ.SPES LLQLKALKPG VIQILGVKTS
mAgp-26257	TDDDQDTEAH	LEIREDGTVV	GAAHR.SPES LLELKALKPG VIQILGVKAS
Hfgf14	CR..QGY..Y	LQMHPDGALD	GTKDDSTNST LFNLI PVGLR VVAIQGVKTG
Mfgf14	CR..QGY..Y	LQMHPDGALD	GTKDDSTNST LFNLI PVGLR VVAIQGVKTG
Hfgf12	SQ..QGY..F	LQMHPDGTID	GTKDENS DYT LFNLI PVGLR VVAIQGVKAS
Mfgf13	SR..QGY..H	LQLQADGTID	GTKDEDSTYT LFNLI PVGLR VVAIQGVQTK
Hfgf5	CRVGIGF..H	LQIYPDGKVN	GSHE.ANMLS VLEIFAVSQG IVGIRGVFSN
Mfgf5	CRVGIGF..H	LQIYPDGKVN	GSHE.ASVLS ILEIFAVSQG IVGIRGVFSN
Hfgf6	CNVGIGF..H	LQVLPDGRIS	GTHE.ENPYS LLEISTVERG VVSLFGVRSA
Mfgf6	CNVGIGF..H	LQVPPDGRIS	GTHE.ENPYS LLEISTVERG VVSLFGVKSA
Hfgf4	CNVGIGF..H	LQALPDGRIG	GAHA.DTRDS LLELSPVERG VVSIFGVASR
Mfgf4	CNVGIGF..H	LQVLPDGRIG	GVHA.DTRDS LLELSPVQRG VVSIFGVASR
Hfgf3	CATK..Y..H	LQLHPSGRVN	GSLE.NSAYS ILEITAVEVG IVAIRGLFSG
Mfgf3	CATK..Y..H	LQLHPSGRVN	GSLE.NSAYS ILEITAVEVG VVAIKGLFSG
Hfgf7	CRTQ..W..Y	LRIDKRGKVK	GTQEMKNYN IMEIRTVAVG IVAIKGVESE
Mfgf7	CRTQ..W..Y	LRIDKRGKVK	GTQEMKNSYN IMEIRTVAVG IVAIKGVESE
Hfgf9	CRT..GF..H	LEIFPNGTIQ	GTRKDH SRFG ILEFISIAVG LVSIRGVDSG
Mfgf9	CRT..GF..H	LEIFPNGTIQ	GTRKDH SRFG ILEFISIAVG LVSIRGVDSG
Hfgf1	CSNG.GH..F	LRILPDGTVD	GTRDRSDQHI QLQLSAESVG EVYIKSTETG
Mfgf1	CSNG.GH..F	LRILPDGTVD	GTRDRSDQHI QLQLSAESAG EVYIKGTETG
Hfgf2	CKNG.GF..F	LRIHPDGRVD	GVREKSDPHI KLQLQAEERG VVSIKGV CAN
Mfgf2	CKNG.GF..F	LRIHPDGRVD	GVREKSDPHV KLQLQAEERG VVSIKGV CAN
cons	cr.g.gf..h	LqihpdG.vd	Gt.e.sspys llel.avevg vv.ikgvkag

	151		200
hAgp-26257	RFLCQRPDGA	LYGSLHFDPE	ACSFRELLLE DGYNVYQSEA HGLPLHLPGN
mAgp-26257	RFLCQQPDGA	LYGSPHFDPE	ACSFRELLLE DGYNVYQSEA HGLPLRLPQK
Hfgf14	LYIAMNGEGY	LYPSELFTPE	.CKFKESVFE NYYVIYSSML YRQQESGRA.
Mfgf14	LYIAMNGEGY	LYPSELFTPE	.CKFKESVFE NYYVIYSSML YRQQESGRA.
Hfgf12	LYVAMNGEGY	LYSSDVFTPE	.CKFKESVFE NYYVIYSSTL YRQQESGRA.
Mfgf13	LYLAMNSEGY	LYTSEHFTPE	.CKFKESVFE NYYVTYSSMI YRQQQSGRG.
Hfgf5	KFLAMSKKGK	LHASAKFTDD	.CKFRERFQE NSYNTYASAI HRTEKTGRE.
Mfgf5	KFLAMSKKGK	LHASAKFTDD	.CKFRERFQE NSYNTYASAI HRTEKTGRE.
Hfgf6	LFVAMNSKGR	LYATPSFQEE	.CKFRETL LP NNYNAYESDL Y.....QG.
Mfgf6	LFIAMNSKGR	LYTTPSFHDE	.CKFRETL LP NNYNAYESDL Y.....RG.
Hfgf4	FFVAMSSKGK	LYGSPFFTDE	.CTFKEILLP NNYNAYESYK Y.....PG.
Mfgf4	FFVAMSSRGK	LFGVPFFTDE	.CKFKEILLP NNYNAYEAYA Y.....PG.
Hfgf3	RYLAMNKRGR	LYASEHYSAE	.CEFVERIHE LGYNTYASRL YRTVSSTPGA
Mfgf3	RYLAMNKRGR	LYASDHYNAE	.CEFVERIHE LGYNTYASRL YRTGSSGPGA
Hfgf7	FYLAMNKEGK	LYAKKECNE	.CNFKELILE NHYNTYASAK WTH..NG...
Mfgf7	YYLAMNKEGK	LYAKKECNE	.CNFKELILE NHYNTYASAK WTH..SG...
Hfgf9	LYLGMNEKGE	LYGSEKLTQE	.CVFREQFEE NWYNTYSSNL YKHVD TG...
Mfgf9	LYLGMNEKGE	LYGSEKLTQE	.CVFREQFEE NWYNTYSSNL YKHVD TG...
Hfgf1	QYLAMDTDGL	LYGSQTPNEE	.CLFLERLEE NHYNTYISK H....AE...
Mfgf1	QYLAMDTGL	LYGSQTPNEE	.CLFLERLEE NHYNTYTSK H....AE...
Hfgf2	RYLAMKEDGR	LLASKCVTDE	.CFFFERLES NNYNTYRSRK Y....T....
Mfgf2	RYLAMKEDGR	LLASKCVTEE	.CFFFERLES NNYNTYRSRK Y....S....

6/10

Fig. 3C

	201				250
hAgp-26257	KSPHRDPAPR	GPA...RFLP	LPGLPPAPPE	PPGILAPQPP	DVGSSDPLSM
mAgp-26257	DSPNQDATSW	GPV...RFLP	MPGLLHEPQD	QAGFLPPEPP	DVGSSDPLSM
Hfgf14	.....W	FLGLNKEGQA	MKG...NRVKK	TKPAAHFLPK	PLEVAMYREP
Mfgf14	.....W	FLGLNKEGQV	MKG...NRVKK	TKPAAHFLPK	PLEVAMYREP
Hfgf12	.....W	FLGLNKEGQI	MKG...NRVKK	TKPSSHFLPK	PIEVCMYREP
Mfgf13	.....W	YLGLNKEGEI	MKG...NHVKK	NKPAAHFLPK	PLKVAMYKEP
Hfgf5	.....W	YVALNKRKGA	KRGCSPRVKP	QHISTHFLPR	FKQSEQPELS
Mfgf5	.....W	YVALNKRKGA	KRGCSPRVKP	QHVSTHFLPR	FKQSEQPELS
Hfgf6	.....T	YIALSKYGRV	KRG...SKVSP	IMTVTHFLPR	I-----
Mfgf6	.....T	YIALSKYGRV	KRG...SKVSP	IMTVTHFLPR	I-----
Hfgf4	.....M	FIALSKNGKT	KKG...NRVSP	TMKVTHFLPR	L-----
Mfgf4	.....M	FMALSKNGRT	KKG...NRVSP	TMKVTHFLPR	L-----
Hfgf3	RRQPSAERLW	YVSVNGKGRP	RRG...FKTRR	TQKSSLFLPR	VLDHRDHEMV
Mfgf3	QRQPGAQRPW	YVSVNGKGRP	RRG...FKTRR	TQKSSLFLPR	VLGHKDHEMV
Hfgf7	.....GEM	FVALNQKGIP	VRG...KKTCK	EQKTAHFLPM	AIT-----
Mfgf7	.....GEM	FVALNQKGIP	VKG...KKTCK	EQKTAHFLPM	AIT-----
Hfgf9	.....RRY	YVALNKDGTP	REG...TRTKR	HQKFTHFLPR	PVDPDKVPEL
Mfgf9	.....RRY	YVALNKDGTP	REG...TRTKR	HQKFTHFLPR	PVDPDKVPEL
Hfgf1	.....KNW	FVGLKKNGSC	KRG...PRTHY	GQKAILFLPL	PVSSD-----
Mfgf1	.....KNW	FVGLKKNGSC	KRG...PRTHY	GQKAILFLPL	PVSSD-----
Hfgf2	.....SW	YVALKRTGQY	KLG...SKTGP	GQKAILFLPM	SAKS-----
Mfgf2	.....SW	YVALKRTGQY	KLG...SKTGP	GQKAILFLPM	SAKS-----
cons	.....w	yvalnk.g.p	krG...nr.kp	tqkathflPr	pv-s-----
	251				300
hAgp-26257	VGPSQGRSPS	YAS-----	-----	-----	-----
mAgp-26257	VEPLQGRSPS	YAS-----	-----	-----	-----
Hfgf14	SLHDVGETVP	KPGVTPSKST	SASAIMNGGK	PVNKSKTT--	-----
Mfgf14	SLHDVGETVP	KAGVTPSKST	SASAIMNGGK	PVNKCKTT--	-----
Hfgf12	SLHEIGEKQG	R.....SRKS	SGTPTMNGGK	VVNQDST---	-----
Mfgf13	SLHDLTEFSR	SGSGTPTKSR	SVSGVLNGGK	SMSHNEST--	-----
Hfgf5	FTVTVPKKN	PPSPIKSKIP	LSAPRKNTNS	VKYRLKFRFG	-----
Mfgf5	FTVTVPKKN	P...PVKPKVP	LSQPRRSPSP	VKYRLKFRFG	-----
Hfgf6	-----	-----	-----	-----	-----
Mfgf6	-----	-----	-----	-----	-----
Hfgf4	-----	-----	-----	-----	-----
Mfgf4	-----	-----	-----	-----	-----
Hfgf3	RQLQSGLPRP	PGKGVQPRRR	RQ.KQSPDNL	EP SHVQASRL	GSQLEASAH-
Mfgf3	RLQSSQPRA	PGEQSQRQR	RQKKQSPGDH	GKMETLSTRA	TPSTQLHTGG
Hfgf7	-----	-----	-----	-----	-----
Mfgf7	-----	-----	-----	-----	-----
Hfgf9	YKDILSQS--	-----	-----	-----	-----
Mfgf9	YKDILSQS--	-----	-----	-----	-----
Hfgf1	-----	-----	-----	-----	-----
Mfgf1	-----	-----	-----	-----	-----
Hfgf2	-----	-----	-----	-----	-----
Mfgf2	-----	-----	-----	-----	-----

Fig. 3D

	301
hAgp-26257	-----
mAgp-26257	-----
Hfgf14	-----
Mfgf14	-----
Hfgf12	-----
Mfgf13	-----
Hfgf5	-----
Mfgf5	-----
Hfgf6	-----
Mfgf6	-----
Hfgf4	-----
Mfgf4	-----
Hfgf3	-----
Mfgf3	LAVA
Hfgf7	-----
Mfgf7	-----
Hfgf9	-----
Mfgf9	-----
Hfgf1	-----
Mfgf1	-----
Hfgf2	-----
Mfgf2	-----
cons	-----

Fig. 4A




Fig. 4B





Fig. 4C



	1	2	3	4	5	6	7	8
A	whole brain	amygdala	caudate nucleus	cerebellum	cerebral cortex	frontal lobe	hippo-campus	medulla oblonggata
B	occipital lobe	putamen	substantia nigra	temporal lobe	thalamus	nucleus accumbens	spinal cord	
C	heart	aorta	skeletal muscle	colon	bladder	uterus	prostate	stomach
D	testis	ovary	pancreas	pituitary gland	adrenal gland	thyroid gland	salivary gland	mammary gland
E	kidney	liver	small intestine	spleen	thymus	peripheral leukocyte	lymph node	bone marrow
F	appendix	lung	trachea	placenta				
G	fetal brain	fetal heart	fetal kidney	fetal liver	fetal spleen	fetal thymus	fetal lung	
H	yeast T-RNA 100 ng	yeast tRNA 100 ng	E. coli rRNA 100 ng	E. coli DNA 100 ng	Poly-(A) 100 ng	human Cot1DNA 100 ng	human DNA 100 ng	human DNA 500 ng



## SEQUENCE LISTING

<110> Thomason, Arlen  
Liu, Benxian

<120> Fibroblast Growth Factor-Like Polypeptides

<130> 99-371-A

<140>

<141>

<150> 09/391,861

<151> 1999-09-07

<160> 41

<170> PatentIn Ver. 2.0

<210> 1

<211> 1190

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (142)..(771)

<400> 1

```

gaggatccag ccgaaagagg agccaggcac tcaggccacc tgagtctact cacctggaca 60
actggaatct ggcaccaatt ctaaaccact cagcttctcc gagctcacac cccggagatc 120
acctgaggac ccgagccatt g atg gac tcg gac gag acc ggg ttc gag cac 171
                        Met Asp Ser Asp Glu Thr Gly Phe Glu His
                        1         5         10
tca gga ctg tgg gtt tct gtg ctg gct ggt ctt ctg ctg gga gcc tgc 219
Ser Gly Leu Trp Val Ser Val Leu Ala Gly Leu Leu Leu Gly Ala Cys
                        15         20         25
cag gca cac ccc atc cct gac tcc agt cct ctc ctg caa ttc ggg ggc 267
Gln Ala His Pro Ile Pro Asp Ser Ser Pro Leu Leu Gln Phe Gly Gly
                        30         35         40
caa gtc cgg cag cgg tac ctc tac aca gat gat gcc cag cag aca gaa 315
Gln Val Arg Gln Arg Tyr Leu Tyr Thr Asp Asp Ala Gln Gln Thr Glu
                        45         50         55
gcc cac ctg gag atc agg gag gat ggg acg gtg ggg ggc gct gct gac 363
Ala His Leu Glu Ile Arg Glu Asp Gly Thr Val Gly Gly Ala Ala Asp
                        60         65         70
cag agc ccc gaa agt ctc ctg cag ctg aaa gcc ttg aag ccg gga gtt 411
Gln Ser Pro Glu Ser Leu Leu Gln Leu Lys Ala Leu Lys Pro Gly Val
                        75         80         85         90

```

att caa atc ttg gga gtc aag aca tcc agg ttc ctg tgc cag cgg cca 459  
 Ile Gln Ile Leu Gly Val Lys Thr Ser Arg Phe Leu Cys Gln Arg Pro  
                   95                  100                  105

gat ggg gcc ctg tat gga tcg ctc cac ttt gac cct gag gcc tgc agc 507  
 Asp Gly Ala Leu Tyr Gly Ser Leu His Phe Asp Pro Glu Ala Cys Ser  
                   110                  115                  120

ttc cgg gag ctg ctt ctt gag gac gga tac aat gtt tac cag tcc gaa 555  
 Phe Arg Glu Leu Leu Leu Glu Asp Gly Tyr Asn Val Tyr Gln Ser Glu  
                   125                  130                  135

gcc cac ggc ctc ccg ctg cac ctg cca ggg aac aag tcc cca cac cgg 603  
 Ala His Gly Leu Pro Leu His Leu Pro Gly Asn Lys Ser Pro His Arg  
                   140                  145                  150

gac cct gca ccc cga gga cca gct cgc ttc ctg cca cta cca ggc ctg 651  
 Asp Pro Ala Pro Arg Gly Pro Ala Arg Phe Leu Pro Leu Pro Gly Leu  
 155                  160                  165                  170

ccc ccc gca ccc ccg gag cca ccc gga atc ctg gcc ccc cag ccc ccc 699  
 Pro Pro Ala Pro Pro Glu Pro Pro Gly Ile Leu Ala Pro Gln Pro Pro  
                   175                  180                  185

gat gtg ggc tcc tcg gac cct ctg agc atg gtg gga cct tcc cag ggc 747  
 Asp Val Gly Ser Ser Asp Pro Leu Ser Met Val Gly Pro Ser Gln Gly  
                   190                  195                  200

cga agc ccc agc tac gct tcc tga agccagaggc tgtttactat gacatctcct 801  
 Arg Ser Pro Ser Tyr Ala Ser  
                   205                  210

ctttatttat taggttattt atcttattta tttttttatt tttcttactt gagataataa 861  
 agagtccag aggaggataa gaatgagcat gtgtgagtgt ctgagggaag acatggcagc 921  
 tgttttgtct cccttggecc ggacaatccc ctctacacct cccctcacgt ggtccgaggg 981  
 tcttggttc ccactgggccc tcaactttttt cttttctttt cttttctttt ttttgagacg 1041  
 gagtctcgct ctgcactcca gccagggcca cagagcgaga ttccatctca aaaaaataaa 1101  
 taaataaata aataaataaa tataaaaata aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1161  
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1190

&lt;210&gt; 2

&lt;211&gt; 209

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 2

Met Asp Ser Asp Glu Thr Gly Phe Glu His Ser Gly Leu Trp Val Ser  
   1                  5                  10                  15

Val Leu Ala Gly Leu Leu Leu Gly Ala Cys Gln Ala His Pro Ile Pro

20	25	30
Asp Ser Ser Pro Leu Leu Gln Phe Gly Gly Gln Val Arg Gln Arg Tyr		
35	40	45
Leu Tyr Thr Asp Asp Ala Gln Gln Thr Glu Ala His Leu Glu Ile Arg		
50	55	60
Glu Asp Gly Thr Val Gly Gly Ala Ala Asp Gln Ser Pro Glu Ser Leu		
65	70	75 80
Leu Gln Leu Lys Ala Leu Lys Pro Gly Val Ile Gln Ile Leu Gly Val		
85	90	95
Lys Thr Ser Arg Phe Leu Cys Gln Arg Pro Asp Gly Ala Leu Tyr Gly		
100	105	110
Ser Leu His Phe Asp Pro Glu Ala Cys Ser Phe Arg Glu Leu Leu Leu		
115	120	125
Glu Asp Gly Tyr Asn Val Tyr Gln Ser Glu Ala His Gly Leu Pro Leu		
130	135	140
His Leu Pro Gly Asn Lys Ser Pro His Arg Asp Pro Ala Pro Arg Gly		
145	150	155 160
Pro Ala Arg Phe Leu Pro Leu Pro Gly Leu Pro Pro Ala Pro Pro Glu		
165	170	175
Pro Pro Gly Ile Leu Ala Pro Gln Pro Pro Asp Val Gly Ser Ser Asp		
180	185	190
Pro Leu Ser Met Val Gly Pro Ser Gln Gly Arg Ser Pro Ser Tyr Ala		
195	200	205

Ser

<210> 3  
 <211> 649  
 <212> DNA  
 <213> Mus musculus

<220>  
 <221> CDS  
 <222> (1) .. (630)

<400> 3														
atg gaa tgg atg aga tct aga gtt ggg acc ctg gga ctg tgg gtc cga	48													
Met Glu Trp Met Arg Ser Arg Val Gly Thr Leu Gly Leu Trp Val Arg														
1 5 10 15														
ctg ctg ctg gct gtc ttc ctg ctg ggg gtc tac caa gca tac ccc atc	96													
Leu Leu Leu Ala Val Phe Leu Leu Gly Val Tyr Gln Ala Tyr Pro Ile														
20 25 30														
cct gac tcc agc ccc ctc ctc cag ttt ggg ggt caa gtc cgg cag agg	144													

Pro Asp Ser Ser Pro Leu Leu Gln Phe Gly Gly Gln Val Arg Gln Arg  
 35 40 45  
 tac ctc tac aca gat gac gac caa gac act gaa gcc cac ctg gag atc 192  
 Tyr Leu Tyr Thr Asp Asp Asp Gln Asp Thr Glu Ala His Leu Glu Ile  
 50 55 60  
 agg gag gat gga aca gtg gta ggc gca gca cac cgc agt cca gaa agt 240  
 Arg Glu Asp Gly Thr Val Val Gly Ala Ala His Arg Ser Pro Glu Ser  
 65 70 75 80  
 ctc ctg gag ctc aaa gcc ttg aag cca ggg gtc att caa atc ctg ggt 288  
 Leu Leu Glu Leu Lys Ala Leu Lys Pro Gly Val Ile Gln Ile Leu Gly  
 85 90 95  
 gtc aaa gcc tct agg ttt ctt tgc caa cag cca gat gga gct ctc tat 336  
 Val Lys Ala Ser Arg Phe Leu Cys Gln Gln Pro Asp Gly Ala Leu Tyr  
 100 105 110  
 gga tcg cct cac ttt gat cct gag gcc tgc agc ttc aga gaa ctg ctg 384  
 Gly Ser Pro His Phe Asp Pro Glu Ala Cys Ser Phe Arg Glu Leu Leu  
 115 120 125  
 ctg gag gac ggt tac aat gtg tac cag tct gaa gcc cat ggc ctg ccc 432  
 Leu Glu Asp Gly Tyr Asn Val Tyr Gln Ser Glu Ala His Gly Leu Pro  
 130 135 140  
 ctg cgt ctg cct cag aag gac tcc cca aac cag gat gca aca tcc tgg 480  
 Leu Arg Leu Pro Gln Lys Asp Ser Pro Asn Gln Asp Ala Thr Ser Trp  
 145 150 155 160  
 gga cct gtg cgc ttc ctg ccc atg cca ggc ctg ctc cac gag ccc caa 528  
 Gly Pro Val Arg Phe Leu Pro Met Pro Gly Leu Leu His Glu Pro Gln  
 165 170 175  
 gac caa gca gga ttc ctg ccc cca gag ccc cca gat gtg ggc tcc tct 576  
 Asp Gln Ala Gly Phe Leu Pro Pro Glu Pro Pro Asp Val Gly Ser Ser  
 180 185 190  
 gac ccc ctg agc atg gta gag cct tta cag ggc cga agc ccc agc tat 624  
 Asp Pro Leu Ser Met Val Glu Pro Leu Gln Gly Arg Ser Pro Ser Tyr  
 195 200 205  
 gcg tcc tgactctttc ctgaatcta 649  
 Ala Ser  
 210  
 <210> 4  
 <211> 210  
 <212> PRT  
 <213> Mus musculus  
 <400> 4  
 Met Glu Trp Met Arg Ser Arg Val Gly Thr Leu Gly Leu Trp Val Arg  
 1 5 10 15

Leu Leu Leu Ala Val Phe Leu Leu Gly Val Tyr Gln Ala Tyr Pro Ile  
                   20                  25                  30  
 Pro Asp Ser Ser Pro Leu Leu Gln Phe Gly Gly Gln Val Arg Gln Arg  
                   35                  40                  45  
 Tyr Leu Tyr Thr Asp Asp Asp Gln Asp Thr Glu Ala His Leu Glu Ile  
                   50                  55                  60  
 Arg Glu Asp Gly Thr Val Val Gly Ala Ala His Arg Ser Pro Glu Ser  
                   65                  70                  75                  80  
 Leu Leu Glu Leu Lys Ala Leu Lys Pro Gly Val Ile Gln Ile Leu Gly  
                   85                  90                  95  
 Val Lys Ala Ser Arg Phe Leu Cys Gln Gln Pro Asp Gly Ala Leu Tyr  
                   100                  105                  110  
 Gly Ser Pro His Phe Asp Pro Glu Ala Cys Ser Phe Arg Glu Leu Leu  
                   115                  120                  125  
 Leu Glu Asp Gly Tyr Asn Val Tyr Gln Ser Glu Ala His Gly Leu Pro  
                   130                  135                  140  
 Leu Arg Leu Pro Gln Lys Asp Ser Pro Asn Gln Asp Ala Thr Ser Trp  
                   145                  150                  155                  160  
 Gly Pro Val Arg Phe Leu Pro Met Pro Gly Leu Leu His Glu Pro Gln  
                   165                  170                  175  
 Asp Gln Ala Gly Phe Leu Pro Pro Glu Pro Pro Asp Val Gly Ser Ser  
                   180                  185                  190  
 Asp Pro Leu Ser Met Val Glu Pro Leu Gln Gly Arg Ser Pro Ser Tyr  
                   195                  200                  205  
 Ala Ser  
                   210

<210> 5  
 <211> 181  
 <212> PRT  
 <213> Homo sapiens

<400> 5  
 His Pro Ile Pro Asp Ser Ser Pro Leu Leu Gln Phe Gly Gly Gln Val  
                   1                  5                  10                  15  
 Arg Gln Arg Tyr Leu Tyr Thr Asp Asp Ala Gln Gln Thr Glu Ala His  
                   20                  25                  30  
 Leu Glu Ile Arg Glu Asp Gly Thr Val Gly Gly Ala Ala Asp Gln Ser  
                   35                  40                  45  
 Pro Glu Ser Leu Leu Gln Leu Lys Ala Leu Lys Pro Gly Val Ile Gln  
                   50                  55                  60

Ile Leu Gly Val Lys Thr Ser Arg Phe Leu Cys Gln Arg Pro Asp Gly  
 65 70 75 80  
 Ala Leu Tyr Gly Ser Leu His Phe Asp Pro Glu Ala Cys Ser Phe Arg  
 85 90 95  
 Glu Leu Leu Leu Glu Asp Gly Tyr Asn Val Tyr Gln Ser Glu Ala His  
 100 105 110  
 Gly Leu Pro Leu His Leu Pro Gly Asn Lys Ser Pro His Arg Asp Pro  
 115 120 125  
 Ala Pro Arg Gly Pro Ala Arg Phe Leu Pro Leu Pro Gly Leu Pro Pro  
 130 135 140  
 Ala Pro Pro Glu Pro Pro Gly Ile Leu Ala Pro Gln Pro Pro Asp Val  
 145 150 155 160  
 Gly Ser Ser Asp Pro Leu Ser Met Val Gly Pro Ser Gln Gly Arg Ser  
 165 170 175  
 Pro Ser Tyr Ala Ser  
 180

<210> 6  
 <211> 181  
 <212> PRT  
 <213> Mus musculus

<400> 6  
 Tyr Pro Ile Pro Asp Ser Ser Pro Leu Leu Gln Phe Gly Gly Gln Val  
 1 5 10 15  
 Arg Gln Arg Tyr Leu Tyr Thr Asp Asp Asp Gln Asp Thr Glu Ala His  
 20 25 30  
 Leu Glu Ile Arg Glu Asp Gly Thr Val Val Gly Ala Ala His Arg Ser  
 35 40 45  
 Pro Glu Ser Leu Leu Glu Leu Lys Ala Leu Lys Pro Gly Val Ile Gln  
 50 55 60  
 Ile Leu Gly Val Lys Ala Ser Arg Phe Leu Cys Gln Gln Pro Asp Gly  
 65 70 75 80  
 Ala Leu Tyr Gly Ser Pro His Phe Asp Pro Glu Ala Cys Ser Phe Arg  
 85 90 95  
 Glu Leu Leu Leu Glu Asp Gly Tyr Asn Val Tyr Gln Ser Glu Ala His  
 100 105 110  
 Gly Leu Pro Leu Arg Leu Pro Gln Lys Asp Ser Pro Asn Gln Asp Ala  
 115 120 125  
 Thr Ser Trp Gly Pro Val Arg Phe Leu Pro Met Pro Gly Leu Leu His



130	135	140
Glu Pro Gln Asp Gln Ala Gly Phe Leu Pro Pro Glu Pro Pro Asp Val		
145	150	155 160
Gly Ser Ser Asp Pro Leu Ser Met Val Glu Pro Leu Gln Gly Arg Ser		
	165	170 175
Pro Ser Tyr Ala Ser		
180		

<210> 7  
 <211> 21  
 <212> DNA  
 <213> Mus musculus

<400> 7  
 tggaatggat gagatctaga g 21

<210> 8  
 <211> 19  
 <212> DNA  
 <213> Mus musculus

<400> 8  
 ctagattcag gaagagtca 19

<210> 9  
 <211> 32  
 <212> DNA  
 <213> Mus musculus

<400> 9  
 cattgcggcc gctcaagatg caaaacgcag tg 32

<210> 10  
 <211> 37  
 <212> DNA  
 <213> Mus musculus

<400> 10  
 ctactaaagc ttccaccatg gaatggatga gatctag 37

<210> 11  
 <211> 33  
 <212> DNA  
 <213> Mus musculus

<400> 11  
 attcatgcgg ccgcggacgc atagctggggg ctt 33

<210> 12  
 <211> 36  
 <212> DNA  
 <213> Homo sapiens

<400> 12  
 ctactaaagc ttccaccatg gactcggacg agaccg . 36

<210> 13  
 <211> 34  
 <212> DNA  
 <213> Homo sapiens

<400> 13  
 attcatgcgg ccgcggaagc gtagctgggg cttc 34

<210> 14  
 <211> 155  
 <212> PRT  
 <213> Homo sapiens

<400> 14  
 Met Ala Glu Gly Glu Ile Thr Thr Phe Thr Ala Leu Thr Glu Lys Phe  
     1                    5                    10                    15  
 Asn Leu Pro Pro Gly Asn Tyr Lys Lys Pro Lys Leu Leu Tyr Cys Ser  
                     20                    25                    30  
 Asn Gly Gly His Phe Leu Arg Ile Leu Pro Asp Gly Thr Val Asp Gly  
                     35                    40                    45  
 Thr Arg Asp Arg Ser Asp Gln His Ile Gln Leu Gln Leu Ser Ala Glu  
                     50                    55                    60  
 Ser Val Gly Glu Val Tyr Ile Lys Ser Thr Glu Thr Gly Gln Tyr Leu  
                     65                    70                    75                    80  
 Ala Met Asp Thr Asp Gly Leu Leu Tyr Gly Ser Gln Thr Pro Asn Glu  
                     85                    90                    95  
 Glu Cys Leu Phe Leu Glu Arg Leu Glu Glu Asn His Tyr Asn Thr Tyr  
                     100                    105                    110  
 Ile Ser Lys Lys His Ala Glu Lys Asn Trp Phe Val Gly Leu Lys Lys  
                     115                    120                    125  
 Asn Gly Ser Cys Lys Arg Gly Pro Arg Thr His Tyr Gly Gln Lys Ala  
                     130                    135                    140  
 Ile Leu Phe Leu Pro Leu Pro Val Ser Ser Asp  
                     145                    150                    155

<210> 15  
 <211> 243

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 15

Met Ala Ala Ala Ile Ala Ser Ser Leu Ile Arg Gln Lys Arg Gln Ala  
 1 5 10 15

Arg Glu Ser Asn Ser Asp Arg Val Ser Ala Ser Lys Arg Arg Ser Ser  
 20 25 30

Pro Ser Lys Asp Gly Arg Ser Leu Cys Glu Arg His Val Leu Gly Val  
 35 40 45

Phe Ser Lys Val Arg Phe Cys Ser Gly Arg Lys Arg Pro Val Arg Arg  
 50 55 60

Arg Pro Glu Pro Gln Leu Lys Gly Ile Val Thr Arg Leu Phe Ser Gln  
 65 70 75 80

Gln Gly Tyr Phe Leu Gln Met His Pro Asp Gly Thr Ile Asp Gly Thr  
 85 90 95

Lys Asp Glu Asn Ser Asp Tyr Thr Leu Phe Asn Leu Ile Pro Val Gly  
 100 105 110

Leu Arg Val Val Ala Ile Gln Gly Val Lys Ala Ser Leu Tyr Val Ala  
 115 120 125

Met Asn Gly Glu Gly Tyr Leu Tyr Ser Ser Asp Val Phe Thr Pro Glu  
 130 135 140

Cys Lys Phe Lys Glu Ser Val Phe Glu Asn Tyr Tyr Val Ile Tyr Ser  
 145 150 155 160

Ser Thr Leu Tyr Arg Gln Gln Glu Ser Gly Arg Ala Trp Phe Leu Gly  
 165 170 175

Leu Asn Lys Glu Gly Gln Ile Met Lys Gly Asn Arg Val Lys Lys Thr  
 180 185 190

Lys Pro Ser Ser His Phe Val Pro Lys Pro Ile Glu Val Cys Met Tyr  
 195 200 205

Arg Glu Pro Ser Leu His Glu Ile Gly Glu Lys Gln Gly Arg Ser Arg  
 210 215 220

Lys Ser Ser Gly Thr Pro Thr Met Asn Gly Gly Lys Val Val Asn Gln  
 225 230 235 240

Asp Ser Thr

&lt;210&gt; 16

&lt;211&gt; 247

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

<400> 16  
 Met Ala Ala Ala Ile Ala Ser Gly Leu Ile Arg Gln Lys Arg Gln Ala  
 1 5 10 15  
 Arg Glu Gln His Trp Asp Arg Pro Ser Ala Ser Arg Arg Arg Ser Ser  
 20 25 30  
 Pro Ser Lys Asn Arg Gly Leu Cys Asn Gly Asn Leu Val Asp Ile Phe  
 35 40 45  
 Ser Lys Val Arg Ile Phe Gly Leu Lys Lys Arg Arg Leu Arg Arg Gln  
 50 55 60  
 Asp Pro Gln Leu Lys Gly Ile Val Thr Arg Leu Tyr Cys Arg Gln Gly  
 65 70 75 80  
 Tyr Tyr Leu Gln Met His Pro Asp Gly Ala Leu Asp Gly Thr Lys Asp  
 85 90 95  
 Asp Ser Thr Asn Ser Thr Leu Phe Asn Leu Ile Pro Val Gly Leu Arg  
 100 105 110  
 Val Val Ala Ile Gln Gly Val Lys Thr Gly Leu Tyr Ile Ala Met Asn  
 115 120 125  
 Gly Glu Gly Tyr Leu Tyr Pro Ser Glu Leu Phe Thr Pro Glu Cys Lys  
 130 135 140  
 Phe Lys Glu Ser Val Phe Glu Asn Tyr Tyr Val Ile Tyr Ser Ser Met  
 145 150 155 160  
 Leu Tyr Arg Gln Gln Glu Ser Gly Arg Ala Trp Phe Leu Gly Leu Asn  
 165 170 175  
 Lys Glu Gly Gln Ala Met Lys Gly Asn Arg Val Lys Lys Thr Lys Pro  
 180 185 190  
 Ala Ala His Phe Leu Pro Lys Pro Leu Glu Val Ala Met Tyr Arg Glu  
 195 200 205  
 Pro Ser Leu His Asp Val Gly Glu Thr Val Pro Lys Pro Gly Val Thr  
 210 215 220  
 Pro Ser Lys Ser Thr Ser Ala Ser Ala Ile Met Asn Gly Gly Lys Pro  
 225 230 235 240  
 Val Asn Lys Ser Lys Thr Thr  
 245

<210> 17  
 <211> 155  
 <212> PRT  
 <213> Homo sapiens

<400> 17

Met Ala Ala Gly Ser Ile Thr Thr Leu Pro Ala Leu Pro Glu Asp Gly  
 1 5 10 15  
 Gly Ser Gly Ala Phe Pro Pro Gly His Phe Lys Asp Pro Lys Arg Leu  
 20 25 30  
 Tyr Cys Lys Asn Gly Gly Phe Phe Leu Arg Ile His Pro Asp Gly Arg  
 35 40 45  
 Val Asp Gly Val Arg Glu Lys Ser Asp Pro His Ile Lys Leu Gln Leu  
 50 55 60  
 Gln Ala Glu Glu Arg Gly Val Val Ser Ile Lys Gly Val Cys Ala Asn  
 65 70 75 80  
 Arg Tyr Leu Ala Met Lys Glu Asp Gly Arg Leu Leu Ala Ser Lys Cys  
 85 90 95  
 Val Thr Asp Glu Cys Phe Phe Phe Glu Arg Leu Glu Ser Asn Asn Tyr  
 100 105 110  
 Asn Thr Tyr Arg Ser Arg Lys Tyr Thr Ser Trp Tyr Val Ala Leu Lys  
 115 120 125  
 Arg Thr Gly Gln Tyr Lys Leu Gly Ser Lys Thr Gly Pro Gly Gln Lys  
 130 135 140  
 Ala Ile Leu Phe Leu Pro Met Ser Ala Lys Ser  
 145 150 155

&lt;210&gt; 18

&lt;211&gt; 239

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 18

Met Gly Leu Ile Trp Leu Leu Leu Leu Ser Leu Leu Glu Pro Gly Trp  
 1 5 10 15  
 Pro Ala Ala Gly Pro Gly Ala Arg Leu Arg Arg Asp Ala Gly Gly Arg  
 20 25 30  
 Gly Gly Val Tyr Glu His Leu Gly Gly Ala Pro Arg Arg Arg Lys Leu  
 35 40 45  
 Tyr Cys Ala Thr Lys Tyr His Leu Gln Leu His Pro Ser Gly Arg Val  
 50 55 60  
 Asn Gly Ser Leu Glu Asn Ser Ala Tyr Ser Ile Leu Glu Ile Thr Ala  
 65 70 75 80  
 Val Glu Val Gly Ile Val Ala Ile Arg Gly Leu Phe Ser Gly Arg Tyr  
 85 90 95  
 Leu Ala Met Asn Lys Arg Gly Arg Leu Tyr Ala Ser Glu His Tyr Ser  
 100 105 110

Ala Glu Cys Glu Phe Val Glu Arg Ile His Glu Leu Gly Tyr Asn Thr  
 115 120 125  
 Tyr Ala Ser Arg Leu Tyr Arg Thr Val Ser Ser Thr Pro Gly Ala Arg  
 130 135 140  
 Arg Gln Pro Ser Ala Glu Arg Leu Trp Tyr Val Ser Val Asn Gly Lys  
 145 150 155 160  
 Gly Arg Pro Arg Arg Gly Phe Lys Thr Arg Arg Thr Gln Lys Ser Ser  
 165 170 175  
 Leu Phe Leu Pro Arg Val Leu Asp His Arg Asp His Glu Met Val Arg  
 180 185 190  
 Gln Leu Gln Ser Gly Leu Pro Arg Pro Pro Gly Lys Gly Val Gln Pro  
 195 200 205  
 Arg Arg Arg Arg Gln Lys Gln Ser Pro Asp Asn Leu Glu Pro Ser His  
 210 215 220  
 Val Gln Ala Ser Arg Leu Gly Ser Gln Leu Glu Ala Ser Ala His  
 225 230 235

<210> 19  
 <211> 206  
 <212> PRT  
 <213> Homo sapiens

<400> 19  
 Met Ser Gly Pro Gly Thr Ala Ala Val Ala Leu Leu Pro Ala Val Leu  
 1 5 10 15  
 Leu Ala Leu Leu Ala Pro Trp Ala Gly Arg Gly Gly Ala Ala Ala Pro  
 20 25 30  
 Thr Ala Pro Asn Gly Thr Leu Glu Ala Glu Leu Glu Arg Arg Trp Glu  
 35 40 45  
 Ser Leu Val Ala Leu Ser Leu Ala Arg Leu Pro Val Ala Ala Gln Pro  
 50 55 60  
 Lys Glu Ala Ala Val Gln Ser Gly Ala Gly Asp Tyr Leu Leu Gly Ile  
 65 70 75 80  
 Lys Arg Leu Arg Arg Leu Tyr Cys Asn Val Gly Ile Gly Phe His Leu  
 85 90 95  
 Gln Ala Leu Pro Asp Gly Arg Ile Gly Gly Ala His Ala Asp Thr Arg  
 100 105 110  
 Asp Ser Leu Leu Glu Leu Ser Pro Val Glu Arg Gly Val Val Ser Ile  
 115 120 125  
 Phe Gly Val Ala Ser Arg Phe Phe Val Ala Met Ser Ser Lys Gly Lys



130		135		140
Leu Tyr Gly Ser Pro Phe Phe Thr Asp Glu Cys Thr Phe Lys Glu Ile				
145		150		155
Leu Leu Pro Asn Asn Tyr Asn Ala Tyr Glu Ser Tyr Lys Tyr Pro Gly				
		165		170
Met Phe Ile Ala Leu Ser Lys Asn Gly Lys Thr Lys Lys Gly Asn Arg				
		180		185
Val Ser Pro Thr Met Lys Val Thr His Phe Leu Pro Arg Leu				
		195		200
				205

&lt;210&gt; 20

&lt;211&gt; 268

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 20

Met Ser Leu Ser Phe Leu Leu Leu Leu Phe Phe Ser His Leu Ile Leu				
1		5		10
Ser Ala Trp Ala His Gly Glu Lys Arg Leu Ala Pro Lys Gly Gln Pro				
		20		25
Gly Pro Ala Ala Thr Asp Arg Asn Pro Ile Gly Ser Ser Ser Arg Gln				
		35		40
Ser Ser Ser Ser Ala Met Ser Ser Ser Ser Ala Ser Ser Ser Pro Ala				
		50		55
Ala Ser Leu Gly Ser Gln Gly Ser Gly Leu Glu Gln Ser Ser Phe Gln				
		65		70
Trp Ser Pro Ser Gly Arg Arg Thr Gly Ser Leu Tyr Cys Arg Val Gly				
		85		90
Ile Gly Phe His Leu Gln Ile Tyr Pro Asp Gly Lys Val Asn Gly Ser				
		100		105
His Glu Ala Asn Met Leu Ser Val Leu Glu Ile Phe Ala Val Ser Gln				
		115		120
Gly Ile Val Gly Ile Arg Gly Val Phe Ser Asn Lys Phe Leu Ala Met				
		130		135
Ser Lys Lys Gly Lys Leu His Ala Ser Ala Lys Phe Thr Asp Asp Cys				
		145		150
Lys Phe Arg Glu Arg Phe Gln Glu Asn Ser Tyr Asn Thr Tyr Ala Ser				
		165		170
Ala Ile His Arg Thr Glu Lys Thr Gly Arg Glu Trp Tyr Val Ala Leu				
		180		185
				190

Asn Lys Arg Gly Lys Ala Lys Arg Gly Cys Ser Pro Arg Val Lys Pro  
 195 200 205

Gln His Ile Ser Thr His Phe Leu Pro Arg Phe Lys Gln Ser Glu Gln  
 210 215 220

Pro Glu Leu Ser Phe Thr Val Thr Val Pro Glu Lys Lys Asn Pro Pro  
 225 230 235 240

Ser Pro Ile Lys Ser Lys Ile Pro Leu Ser Ala Pro Arg Lys Asn Thr  
 245 250 255

Asn Ser Val Lys Tyr Arg Leu Lys Phe Arg Phe Gly  
 260 265

<210> 21

<211> 208

<212> PRT

<213> Homo sapiens

<400> 21

Met Ala Leu Gly Gln Lys Leu Phe Ile Thr Met Ser Arg Gly Ala Gly  
 1 5 10 15

Arg Leu Gln Gly Thr Leu Trp Ala Leu Val Phe Leu Gly Ile Leu Val  
 20 25 30

Gly Met Val Val Pro Ser Pro Ala Gly Thr Arg Ala Asn Asn Thr Leu  
 35 40 45

Leu Asp Ser Arg Gly Trp Gly Thr Leu Leu Ser Arg Ser Arg Ala Gly  
 50 55 60

Leu Ala Gly Glu Ile Ala Gly Val Asn Trp Glu Ser Gly Tyr Leu Val  
 65 70 75 80

Gly Ile Lys Arg Gln Arg Arg Leu Tyr Cys Asn Val Gly Ile Gly Phe  
 85 90 95

His Leu Gln Val Leu Pro Asp Gly Arg Ile Ser Gly Thr His Glu Glu  
 100 105 110

Asn Pro Tyr Ser Leu Leu Glu Ile Ser Thr Val Glu Arg Gly Val Val  
 115 120 125

Ser Leu Phe Gly Val Arg Ser Ala Leu Phe Val Ala Met Asn Ser Lys  
 130 135 140

Gly Arg Leu Tyr Ala Thr Pro Ser Phe Gln Glu Glu Cys Lys Phe Arg  
 145 150 155 160

Glu Thr Leu Leu Pro Asn Asn Tyr Asn Ala Tyr Glu Ser Asp Leu Tyr  
 165 170 175

Gln Gly Thr Tyr Ile Ala Leu Ser Lys Tyr Gly Arg Val Lys Arg Gly  
 180 185 190

Ser Lys Val Ser Pro Ile Met Thr Val Thr His Phe Leu Pro Arg Ile  
 195 200 205

<210> 22  
 <211> 194  
 <212> PRT  
 <213> Homo sapiens

<400> 22  
 Met His Lys Trp Ile Leu Thr Trp Ile Leu Pro Thr Leu Leu Tyr Arg  
 1 5 10 15  
 Ser Cys Phe His Ile Ile Cys Leu Val Gly Thr Ile Ser Leu Ala Cys  
 20 25 30  
 Asn Asp Met Thr Pro Glu Gln Met Ala Thr Asn Val Asn Cys Ser Ser  
 35 40 45  
 Pro Glu Arg His Thr Arg Ser Tyr Asp Tyr Met Glu Gly Gly Asp Ile  
 50 55 60  
 Arg Val Arg Arg Leu Phe Cys Arg Thr Gln Trp Tyr Leu Arg Ile Asp  
 65 70 75 80  
 Lys Arg Gly Lys Val Lys Gly Thr Gln Glu Met Lys Asn Asn Tyr Asn  
 85 90 95  
 Ile Met Glu Ile Arg Thr Val Ala Val Gly Ile Val Ala Ile Lys Gly  
 100 105 110  
 Val Glu Ser Glu Phe Tyr Leu Ala Met Asn Lys Glu Gly Lys Leu Tyr  
 115 120 125  
 Ala Lys Lys Glu Cys Asn Glu Asp Cys Asn Phe Lys Glu Leu Ile Leu  
 130 135 140  
 Glu Asn His Tyr Asn Thr Tyr Ala Ser Ala Lys Trp Thr His Asn Gly  
 145 150 155 160  
 Gly Glu Met Phe Val Ala Leu Asn Gln Lys Gly Ile Pro Val Arg Gly  
 165 170 175  
 Lys Lys Thr Lys Lys Glu Gln Lys Thr Ala His Phe Leu Pro Met Ala  
 180 185 190

Ile Thr

<210> 23  
 <211> 208  
 <212> PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 23

Met Ala Pro Leu Gly Glu Val Gly Asn Tyr Phe Gly Val Gln Asp Ala  
 1 5 10 15

Val Pro Phe Gly Asn Val Pro Val Leu Pro Val Asp Ser Pro Val Leu  
 20 25 30

Leu Ser Asp His Leu Gly Gln Ser Glu Ala Gly Gly Leu Pro Arg Gly  
 35 40 45

Pro Ala Val Thr Asp Leu Asp His Leu Lys Gly Ile Leu Arg Arg Arg  
 50 55 60

Gln Leu Tyr Cys Arg Thr Gly Phe His Leu Glu Ile Phe Pro Asn Gly  
 65 70 75 80

Thr Ile Gln Gly Thr Arg Lys Asp His Ser Arg Phe Gly Ile Leu Glu  
 85 90 95

Phe Ile Ser Ile Ala Val Gly Leu Val Ser Ile Arg Gly Val Asp Ser  
 100 105 110

Gly Leu Tyr Leu Gly Met Asn Glu Lys Gly Glu Leu Tyr Gly Ser Glu  
 115 120 125

Lys Leu Thr Gln Glu Cys Val Phe Arg Glu Gln Phe Glu Glu Asn Trp  
 130 135 140

Tyr Asn Thr Tyr Ser Ser Asn Leu Tyr Lys His Val Asp Thr Gly Arg  
 145 150 155 160

Arg Tyr Tyr Val Ala Leu Asn Lys Asp Gly Thr Pro Arg Glu Gly Thr  
 165 170 175

Arg Thr Lys Arg His Gln Lys Phe Thr His Phe Leu Pro Arg Pro Val  
 180 185 190

Asp Pro Asp Lys Val Pro Glu Leu Tyr Lys Asp Ile Leu Ser Gln Ser  
 195 200 205

&lt;210&gt; 24

&lt;211&gt; 155

&lt;212&gt; PRT

&lt;213&gt; Mus musculus

&lt;400&gt; 24

Met Ala Glu Gly Glu Ile Thr Thr Phe Ala Ala Leu Thr Glu Arg Phe  
 1 5 10 15

Asn Leu Pro Leu Gly Asn Tyr Lys Lys Pro Lys Leu Leu Tyr Cys Ser  
 20 25 30

Asn Gly Gly His Phe Leu Arg Ile Leu Pro Asp Gly Thr Val Asp Gly  
           35                          40                          45  
 Thr Arg Asp Arg Ser Asp Gln His Ile Gln Leu Gln Leu Ser Ala Glu  
           50                          55                          60  
 Ser Ala Gly Glu Val Tyr Ile Lys Gly Thr Glu Thr Gly Gln Tyr Leu  
           65                          70                          75                          80  
 Ala Met Asp Thr Glu Gly Leu Leu Tyr Gly Ser Gln Thr Pro Asn Glu  
                           85                          90                          95  
 Glu Cys Leu Phe Leu Glu Arg Leu Glu Glu Asn His Tyr Asn Thr Tyr  
                           100                          105                          110  
 Thr Ser Lys Lys His Ala Glu Lys Asn Trp Phe Val Gly Leu Lys Lys  
           115                          120                          125  
 Asn Gly Ser Cys Lys Arg Gly Pro Arg Thr His Tyr Gly Gln Lys Ala  
           130                          135                          140  
 Ile Leu Phe Leu Pro Leu Pro Val Ser Ser Asp  
   145                          150                          155

<210> 25  
 <211> 245  
 <212> PRT  
 <213> Mus musculus

<400> 25  
 Met Thr Ala Ala Ile Ala Ser Ser Leu Ile Arg Gln Lys Arg Gln Ala  
   1                          5                          10                          15  
 Arg Glu Arg Glu Lys Ser Asn Ala Cys Lys Cys Val Ser Ser Pro Ser  
           20                          25                          30  
 Lys Gly Lys Thr Ser Cys Asp Lys Asn Lys Leu Asn Val Phe Ser Arg  
           35                          40                          45  
 Val Lys Leu Phe Gly Ser Lys Lys Arg Arg Arg Arg Arg Pro Glu Pro  
           50                          55                          60  
 Gln Leu Lys Gly Ile Val Thr Lys Leu Tyr Ser Arg Gln Gly Tyr His  
           65                          70                          75                          80  
 Leu Gln Leu Gln Ala Asp Gly Thr Ile Asp Gly Thr Lys Asp Glu Asp  
                           85                          90                          95  
 Ser Thr Tyr Thr Leu Phe Asn Leu Ile Pro Val Gly Leu Arg Val Val  
           100                          105                          110  
 Ala Ile Gln Gly Val Gln Thr Lys Leu Tyr Leu Ala Met Asn Ser Glu  
           115                          120                          125  
 Gly Tyr Leu Tyr Thr Ser Glu His Phe Thr Pro Glu Cys Lys Phe Lys

130                      135                      140  
 Glu Ser Val Phe Glu Asn Tyr Tyr Val Thr Tyr Ser Ser Met Ile Tyr  
 145                      150                      155                      160  
 Arg Gln Gln Gln Ser Gly Arg Gly Trp Tyr Leu Gly Leu Asn Lys Glu  
 165                      170                      175  
 Gly Glu Ile Met Lys Gly Asn His Val Lys Lys Asn Lys Pro Ala Ala  
 180                      185                      190  
 His Phe Leu Pro Lys Pro Leu Lys Val Ala Met Tyr Lys Glu Pro Ser  
 195                      200                      205  
 Leu His Asp Leu Thr Glu Phe Ser Arg Ser Gly Ser Gly Thr Pro Thr  
 210                      215                      220  
 Lys Ser Arg Ser Val Ser Gly Val Leu Asn Gly Gly Lys Ser Met Ser  
 225                      230                      235                      240  
 His Asn Glu Ser Thr  
 245

<210> 26  
 <211> 247  
 <212> PRT  
 <213> Mus musculus

<400> 26  
 Met Ala Ala Ala Ile Ala Ser Gly Leu Ile Arg Gln Lys Arg Gln Ala  
 1                      5                      10                      15  
 Arg Glu Gln His Trp Asp Arg Pro Ser Ala Ser Arg Arg Arg Ser Ser  
 20                      25                      30  
 Pro Ser Lys Asn Arg Gly Leu Phe Asn Gly Asn Leu Val Asp Ile Phe  
 35                      40                      45  
 Ser Lys Val Arg Ile Phe Gly Leu Lys Lys Arg Arg Leu Arg Arg Gln  
 50                      55                      60  
 Asp Pro Gln Leu Lys Gly Ile Val Thr Arg Leu Tyr Cys Arg Gln Gly  
 65                      70                      75                      80  
 Tyr Tyr Leu Gln Met His Pro Asp Gly Ala Leu Asp Gly Thr Lys Asp  
 85                      90                      95  
 Asp Ser Thr Asn Ser Thr Leu Phe Asn Leu Ile Pro Val Gly Leu Arg  
 100                      105                      110  
 Val Val Ala Ile Gln Gly Val Lys Thr Gly Leu Tyr Ile Ala Met Asn  
 115                      120                      125  
 Gly Glu Gly Tyr Leu Tyr Pro Ser Glu Leu Phe Thr Pro Glu Cys Lys  
 130                      135                      140



Phe Lys Glu Ser Val Phe Glu Asn Tyr Tyr Val Ile Tyr Ser Ser Met  
 145 150 155 160  
 Leu Tyr Arg Gln Gln Glu Ser Gly Arg Ala Trp Phe Leu Gly Leu Asn  
 165 170 175  
 Lys Glu Gly Gln Val Met Lys Gly Asn Arg Val Lys Lys Thr Lys Pro  
 180 185 190  
 Ala Ala His Phe Leu Pro Lys Pro Leu Glu Val Ala Met Tyr Arg Glu  
 195 200 205  
 Pro Ser Leu His Asp Val Gly Glu Thr Val Pro Lys Ala Gly Val Thr  
 210 215 220  
 Pro Ser Lys Ser Thr Ser Ala Ser Ala Ile Met Asn Gly Gly Lys Pro  
 225 230 235 240  
 Val Asn Lys Cys Lys Thr Thr  
 245

<210> 27  
 <211> 154  
 <212> PRT  
 <213> Mus musculus

<400> 27  
 Met Ala Ala Ser Gly Ile Thr Ser Leu Pro Ala Leu Pro Glu Asp Gly  
 1 5 10 15  
 Gly Ala Ala Phe Pro Pro Gly His Phe Lys Asp Pro Lys Arg Leu Tyr  
 20 25 30  
 Cys Lys Asn Gly Gly Phe Phe Leu Arg Ile His Pro Asp Gly Arg Val  
 35 40 45  
 Asp Gly Val Arg Glu Lys Ser Asp Pro His Val Lys Leu Gln Leu Gln  
 50 55 60  
 Ala Glu Glu Arg Gly Val Val Ser Ile Lys Gly Val Cys Ala Asn Arg  
 65 70 75 80  
 Tyr Leu Ala Met Lys Glu Asp Gly Arg Leu Leu Ala Ser Lys Cys Val  
 85 90 95  
 Thr Glu Glu Cys Phe Phe Phe Glu Arg Leu Glu Ser Asn Asn Tyr Asn  
 100 105 110  
 Thr Tyr Arg Ser Arg Lys Tyr Ser Ser Trp Tyr Val Ala Leu Lys Arg  
 115 120 125  
 Thr Gly Gln Tyr Lys Leu Gly Ser Lys Thr Gly Pro Gly Gln Lys Ala  
 130 135 140  
 Ile Leu Phe Leu Pro Met Ser Ala Lys Ser  
 145 150

&lt;210&gt; 28

&lt;211&gt; 245

&lt;212&gt; PRT

&lt;213&gt; Mus musculus.

&lt;400&gt; 28

Met Gly Leu Ile Trp Leu Leu Leu Leu Ser Leu Leu Glu Pro Ser Trp  
 1 5 10 15  
 Pro Thr Thr Gly Pro Gly Thr Arg Leu Arg Arg Asp Ala Gly Gly Arg  
 20 25 30  
 Gly Gly Val Tyr Glu His Leu Gly Gly Ala Pro Arg Arg Arg Lys Leu  
 35 40 45  
 Tyr Cys Ala Thr Lys Tyr His Leu Gln Leu His Pro Ser Gly Arg Val  
 50 55 60  
 Asn Gly Ser Leu Glu Asn Ser Ala Tyr Ser Ile Leu Glu Ile Thr Ala  
 65 70 75 80  
 Val Glu Val Gly Val Val Ala Ile Lys Gly Leu Phe Ser Gly Arg Tyr  
 85 90 95  
 Leu Ala Met Asn Lys Arg Gly Arg Leu Tyr Ala Ser Asp His Tyr Asn  
 100 105 110  
 Ala Glu Cys Glu Phe Val Glu Arg Ile His Glu Leu Gly Tyr Asn Thr  
 115 120 125  
 Tyr Ala Ser Arg Leu Tyr Arg Thr Gly Ser Ser Gly Pro Gly Ala Gln  
 130 135 140  
 Arg Gln Pro Gly Ala Gln Arg Pro Trp Tyr Val Ser Val Asn Gly Lys  
 145 150 155 160  
 Gly Arg Pro Arg Arg Gly Phe Lys Thr Arg Arg Thr Gln Lys Ser Ser  
 165 170 175  
 Leu Phe Leu Pro Arg Val Leu Gly His Lys Asp His Glu Met Val Arg  
 180 185 190  
 Leu Leu Gln Ser Ser Gln Pro Arg Ala Pro Gly Glu Gly Ser Gln Pro  
 195 200 205  
 Arg Gln Arg Arg Gln Lys Lys Gln Ser Pro Gly Asp His Gly Lys Met  
 210 215 220  
 Glu Thr Leu Ser Thr Arg Ala Thr Pro Ser Thr Gln Leu His Thr Gly  
 225 230 235 240  
 Gly Leu Ala Val Ala  
 245

<210> 29  
 <211> 202  
 <212> PRT  
 <213> Mus musculus

<400> 29  
 Met Ala Lys Arg Gly Pro Thr Thr Gly Thr Leu Leu Pro Arg Val Leu  
 1 5 10 15  
 Leu Ala Leu Val Val Ala Leu Ala Asp Arg Gly Thr Ala Ala Pro Asn  
 20 25 30  
 Gly Thr Arg His Ala Glu Leu Gly His Gly Trp Asp Gly Leu Val Ala  
 35 40 45  
 Arg Ser Leu Ala Arg Leu Pro Val Ala Ala Gln Pro Pro Gln Ala Ala  
 50 55 60  
 Val Arg Ser Gly Ala Gly Asp Tyr Leu Leu Gly Leu Lys Arg Leu Arg  
 65 70 75 80  
 Arg Leu Tyr Cys Asn Val Gly Ile Gly Phe His Leu Gln Val Leu Pro  
 85 90 95  
 Asp Gly Arg Ile Gly Gly Val His Ala Asp Thr Arg Asp Ser Leu Leu  
 100 105 110  
 Glu Leu Ser Pro Val Gln Arg Gly Val Val Ser Ile Phe Gly Val Ala  
 115 120 125  
 Ser Arg Phe Phe Val Ala Met Ser Ser Arg Gly Lys Leu Phe Gly Val  
 130 135 140  
 Pro Phe Phe Thr Asp Glu Cys Lys Phe Lys Glu Ile Leu Leu Pro Asn  
 145 150 155 160  
 Asn Tyr Asn Ala Tyr Glu Ala Tyr Ala Tyr Pro Gly Met Phe Met Ala  
 165 170 175  
 Leu Ser Lys Asn Gly Arg Thr Lys Lys Gly Asn Arg Val Ser Pro Thr  
 180 185 190  
 Met Lys Val Thr His Phe Leu Pro Arg Leu  
 195 200

<210> 30  
 <211> 264  
 <212> PRT  
 <213> Mus musculus

<400> 30  
 Met Ser Leu Ser Leu Leu Phe Leu Ile Phe Cys Ser His Leu Ile His  
 1 5 10 15  
 Ser Ala Trp Ala His Gly Glu Lys Arg Leu Thr Pro Glu Gly Gln Pro  
 20 25 30

Ala Pro Pro Arg Asn Pro Gly Asp Ser Ser Gly Ser Arg Gly Arg Ser  
           35                          40                          45  
 Ser Ala Thr Phe Ser Ser Ser Ser Ala Ser Ser Pro Val Ala Ala Ser  
       50                                  55                          60  
 Pro Gly Ser Gln Gly Ser Gly Ser Glu His Ser Ser Phe Gln Trp Ser  
       65                                  70                          75                          80  
 Pro Ser Gly Arg Arg Thr Gly Ser Leu Tyr Cys Arg Val Gly Ile Gly  
                                   85                          90                          95  
 Phe His Leu Gln Ile Tyr Pro Asp Gly Lys Val Asn Gly Ser His Glu  
                           100                          105                          110  
 Ala Ser Val Leu Ser Ile Leu Glu Ile Phe Ala Val Ser Gln Gly Ile  
           115                          120                          125  
 Val Gly Ile Arg Gly Val Phe Ser Asn Lys Phe Leu Ala Met Ser Lys  
       130                          135                          140  
 Lys Gly Lys Leu His Ala Ser Ala Lys Phe Thr Asp Asp Cys Lys Phe  
       145                          150                          155                          160  
 Arg Glu Arg Phe Gln Glu Asn Ser Tyr Asn Thr Tyr Ala Ser Ala Ile  
                           165                          170                          175  
 His Arg Thr Glu Lys Thr Gly Arg Glu Trp Tyr Val Ala Leu Asn Lys  
                           180                          185                          190  
 Arg Gly Lys Ala Lys Arg Gly Cys Ser Pro Arg Val Lys Pro Gln His  
           195                          200                          205  
 Val Ser Thr His Phe Leu Pro Arg Phe Lys Gln Ser Glu Gln Pro Glu  
       210                          215                          220  
 Leu Ser Phe Thr Val Thr Val Pro Glu Lys Lys Lys Pro Pro Val Lys  
       225                          230                          235                          240  
 Pro Lys Val Pro Leu Ser Gln Pro Arg Arg Ser Pro Ser Pro Val Lys  
                           245                          250                          255  
 Tyr Arg Leu Lys Phe Arg Phe Gly  
           260

&lt;210&gt; 31

&lt;211&gt; 208

&lt;212&gt; PRT

&lt;213&gt; Mus musculus

&lt;400&gt; 31

Met Ala Leu Gly Gln Arg Leu Phe Ile Thr Met Ser Arg Gly Ala Gly  
       1                          5                          10                          15

Arg Val Gln Gly Thr Leu Gln Ala Leu Val Phe Leu Gly Val Leu Val

	20		25		30
Gly Met Val Val Pro Ser Pro Ala Gly Ala Arg Ala Asn Gly Thr Leu	35	40	45		
Leu Asp Ser Arg Gly Trp Gly Thr Leu Leu Ser Arg Ser Arg Ala Gly	50	55	60		
Leu Ala Gly Glu Ile Ser Gly Val Asn Trp Glu Ser Gly Tyr Leu Val	65	70	75	80	
Gly Ile Lys Arg Gln Arg Arg Leu Tyr Cys Asn Val Gly Ile Gly Phe	85	90	95		
His Leu Gln Val Pro Pro Asp Gly Arg Ile Ser Gly Thr His Glu Glu	100	105	110		
Asn Pro Tyr Ser Leu Leu Glu Ile Ser Thr Val Glu Arg Gly Val Val	115	120	125		
Ser Leu Phe Gly Val Lys Ser Ala Leu Phe Ile Ala Met Asn Ser Lys	130	135	140		
Gly Arg Leu Tyr Thr Thr Pro Ser Phe His Asp Glu Cys Lys Phe Arg	145	150	155	160	
Glu Thr Leu Leu Pro Asn Asn Tyr Asn Ala Tyr Glu Ser Asp Leu Tyr	165	170	175		
Arg Gly Thr Tyr Ile Ala Leu Ser Lys Tyr Gly Arg Val Lys Arg Gly	180	185	190		
Ser Lys Val Ser Pro Ile Met Thr Val Thr His Phe Leu Pro Arg Ile	195	200	205		

<210> 32  
 <211> 194  
 <212> PRT  
 <213> Mus musculus

<400> 32  
 Met Arg Lys Trp Ile Leu Thr Arg Ile Leu Pro Thr Leu Leu Tyr Arg  
 1 5 10 15  
 Ser Cys Phe His Leu Val Cys Leu Val Gly Thr Ile Ser Leu Ala Cys  
 20 25 30  
 Asn Asp Met Ser Pro Glu Gln Thr Ala Thr Ser Val Asn Cys Ser Ser  
 35 40 45  
 Pro Glu Arg His Thr Arg Ser Tyr Asp Tyr Met Glu Gly Gly Asp Ile  
 50 55 60

Arg Val Arg Arg Leu Phe Cys Arg Thr Gln Trp Tyr Leu Arg Ile Asp  
 65 70 75 80  
 Lys Arg Gly Lys Val Lys Gly Thr Gln Glu Met Lys Asn Ser Tyr Asn  
 85 90 95  
 Ile Met Glu Ile Arg Thr Val Ala Val Gly Ile Val Ala Ile Lys Gly  
 100 105 110  
 Val Glu Ser Glu Tyr Tyr Leu Ala Met Asn Lys Glu Gly Lys Leu Tyr  
 115 120 125  
 Ala Lys Lys Glu Cys Asn Glu Asp Cys Asn Phe Lys Glu Leu Ile Leu  
 130 135 140  
 Glu Asn His Tyr Asn Thr Tyr Ala Ser Ala Lys Trp Thr His Ser Gly  
 145 150 155 160  
 Gly Glu Met Phe Val Ala Leu Asn Gln Lys Gly Ile Pro Val Lys Gly  
 165 170 175  
 Lys Lys Thr Lys Lys Glu Gln Lys Thr Ala His Phe Leu Pro Met Ala  
 180 185 190  
 Ile Thr

<210> 33  
 <211> 208  
 <212> PRT  
 <213> Mus musculus

<400> 33  
 Met Ala Pro Leu Gly Glu Val Gly Ser Tyr Phe Gly Val Gln Asp Ala  
 1 5 10 15  
 Val Pro Phe Gly Asn Val Pro Val Leu Pro Val Asp Ser Pro Val Leu  
 20 25 30  
 Leu Asn Asp His Leu Gly Gln Ser Glu Ala Gly Gly Leu Pro Arg Gly  
 35 40 45  
 Pro Ala Val Thr Asp Leu Asp His Leu Lys Gly Ile Leu Arg Arg Arg  
 50 55 60  
 Gln Leu Tyr Cys Arg Thr Gly Phe His Leu Glu Ile Phe Pro Asn Gly  
 65 70 75 80  
 Thr Ile Gln Gly Thr Arg Lys Asp His Ser Arg Phe Gly Ile Leu Glu  
 85 90 95  
 Phe Ile Ser Ile Ala Val Gly Leu Val Ser Ile Arg Gly Val Asp Ser  
 100 105 110  
 Gly Leu Tyr Leu Gly Met Asn Glu Lys Gly Glu Leu Tyr Gly Ser Glu  
 115 120 125



Lys Leu Thr Gln Glu Cys Val Phe Arg Glu Gln Phe Glu Glu Asn Trp  
 130 135 140  
 Tyr Asn Thr Tyr Ser Ser Asn Leu Tyr Lys His Val Asp Thr Gly Arg  
 145 150 155 160  
 Arg Tyr Tyr Val Ala Leu Asn Lys Asp Gly Thr Pro Arg Glu Gly Thr  
 165 170 175  
 Arg Thr Lys Arg His Gln Lys Phe Thr His Phe Leu Pro Arg Pro Val  
 180 185 190  
 Asp Pro Asp Lys Val Pro Glu Leu Tyr Lys Asp Ile Leu Ser Gln Ser  
 195 200 205

<210> 34  
 <211> 33  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Random primer  
 with Not I restriction site for first strand cDNA  
 synthesis

<220>  
 <221> misc\_feature  
 <222> (25)..(33)  
 <223> "N" can be A, G, C, or T

<400> 34  
 ggaaggaaaa aagcgccgc aacannnnnn nnn

33

<210> 35  
 <211> 23  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer for  
 first strand cDNA synthesis

<400> 35  
 aatccgatgc ccacgttgca gta

23

<210> 36  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: PCR primer for  
 amplification of cDNA

<400> 36  
aaaatcttag accgacgact gtgttt

26

<210> 37  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: PCR primer for  
amplification of cDNA

<400> 37  
gagtctccgc agccttttga gg

22

<210> 38  
<211> 34  
<212> DNA  
<213> Homo sapiens

<400> 38  
actggcggcc gcaggcatca tcccagttga ggag

34

<210> 39  
<211> 36  
<212> DNA  
<213> Homo sapiens

<400> 39  
actggtcact cgagggtacc ttagctagcc cccggg

36

<210> 40  
<211> 29  
<212> PRT  
<213> Mus musculus

<400> 40  
Met Glu Trp Met Arg Ser Arg Val Gly Thr Leu Gly Leu Trp Val Arg  
1 5 10 15

Leu Leu Leu Ala Val Phe Leu Leu Gly Val Tyr Gln Ala  
20 25

<210> 41  
<211> 28  
<212> PRT  
<213> Homo sapiens

<400> 41  
Met Asp Ser Asp Glu Thr Gly Phe Glu His Ser Gly Leu Trp Val Ser  
1 5 10 15

Val Leu Ala Gly Leu Leu Leu Gly Ala Cys Gln Ala  
20 25

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
15 March 2001 (15.03.2001)

PCT

(10) International Publication Number  
**WO 01/18172 A3**

(51) International Patent Classification<sup>7</sup>: **C12N 15/19**,  
C07K 14/52, G01N 33/50, 33/68, C07K 16/24, G01N  
33/53, A61K 38/19, C07K 19/00, C12Q 1/68, A01K  
67/027, C12N 15/11

2528 Rutland Place, Thousand Oaks, CA 91362 (US).  
**DANILENKO, Dimitri Michael** [US/US]; 750 Rolling  
Oaks Drive, Thousand Oaks, CA 91361 (US).

(21) International Application Number: PCT/US00/24373

(74) Agent: **NOONAN, Kevin, E.**; McDonnell Boehnen Hul-  
bert & Berghoff, 300 South Wacker Drive, Suite 3200,  
Chicago, IL 60606 (US).

(22) International Filing Date:  
5 September 2000 (05.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/391,861 7 September 1999 (07.09.1999) US  
09/644,052 23 August 2000 (23.08.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,  
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(63) Related by continuation (CON) or continuation-in-part  
(CIP) to earlier applications:

US 09/391,861 (CON)  
Filed on 7 September 1999 (07.09.1999)  
US 09/644,052 (CON)  
Filed on 23 August 2000 (23.08.2000)

Published:

-- with international search report

(71) Applicant (*for all designated States except US*): **AMGEN**,  
INC. [US/US]; One Amgen Center Drive, Thousand Oaks,  
CA 91320 (US).

(88) Date of publication of the international search report:  
2 May 2002

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **THOMASON**,  
**Arlen, Read** [US/US]; 2298 Watertown Court, Thou-  
sand Oaks, CA 91360 (US). **LIU, Benxian** [CN/US];

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*



**WO 01/18172 A3**

(54) Title: FIBROBLAST GROWTH FACTOR-LIKE POLYPEPTIDES

(57) Abstract: The present invention provides Fibroblast Growth Factor-like (FGF-like) polypeptides and nucleic acid molecules encoding the same. The invention also provides vectors, host cells, antibodies and methods for producing FGF-like polypeptides. Also provided for are methods for the diagnosis and treatment of diseases associated with FGF-like polypeptides.

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 00/24373

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/19 C07K14/52 G01N33/50 G01N33/68 C07K16/24  
 G01N33/53 A61K38/19 C07K19/00 C12Q1/68 A01K67/027  
 C12N15/11

According to International Patent Classification (IPC) or to both national classification and IPC.

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K G01N A61K C12Q A01K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, SEQUENCE SEARCH

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	WO 01 32678 A (SMITHKLINE BEECHAM CORP (US); AGARWAL; BARONE; FEILD; KABNICK MCQUENEY) 10 May 2001 (2001-05-10) SEQ ID NO:1,2 the whole document ---	1-26, 29-32, 35,37-41
E	WO 01 38357 A (MILLENNIUM PHARMACEUTICALS, INC. (US); KHODADOUST MEHRAN MOHAMAD) 31 May 2001 (2001-05-31) the whole document figure 1 --- -/--	1-29, 31-33, 35,37-41

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

25 January 2002

Date of mailing of the international search report

13/02/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Macchia, G

## INTERNATIONAL SEARCH REPORT

International Application No

PC1/US 00/24373

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
E	<p>WO 01 36640 A (CHIRON CORP (US); KYOTO UNIVERSITY (JP); ITOH N.; KAVANAUGH W. MICHAEL) 25 May 2001 (2001-05-25)</p> <p>the whole document figure 5</p> <p>---</p>	<p>1-12, 14-24, 26-29, 31-33, 35-39, 41</p>
E	<p>WO 01 18209 A (CURAGEN CORP (US) SHIMKETS VERNET BURGESS FERNANDES TAUPIER QUINN ET A) 15 March 2001 (2001-03-15)</p> <p>the whole document page 7</p> <p>---</p>	<p>1-12, 14-24, 26-32, 35, 37-41</p>
E	<p>WO 01 49849 A (ZYMOGENETICS INC. (US); CONKLIN DARRELL C.) 12 July 2001 (2001-07-12)</p> <p>SEQ ID NO:1.2 the whole document</p> <p>---</p>	<p>1-12, 14-23, 26, 29-33, 35, 37-41</p>
E	<p>WO 00 54813 A (CHIRON CORP (US) MANNING DWARKI RENDAHL ZHOU MCGEE LAU FLANNERY MILLER) 21 September 2000 (2000-09-21) page 8; figure 30 the whole document</p> <p>---</p>	<p>1-6, 9, 11, 13-17</p>
X	<p>DATABASE EMBL 'Online! ID: AB006136; Accession number: AB006136, 27 August 1997 (1997-08-27) KODA Y.: "Homo sapiens gene for alpha 1,2-fucosyltransferase, 5' flanking region and partial cds." XP002165898 abstract</p> <p>---</p>	<p>1-5, 7, 11</p>
X	<p>DATABASE EMBL 'Online! ID: AQ175436; Accession number: AQ175436, 21 September 1998 (1998-09-21) MAKAHAIRAS G.G. ET AL.: "HS_3207_A1_H09_MR CIT approved genomic sperm library D Homo sapiens genomic clone plate=3207 col=17 row=0, genomic survey sequence" XP002188227 abstract</p> <p>---</p>	<p>1-5, 7, 11</p>
	<p>---</p> <p>-/--</p>	



# INTERNATIONAL SEARCH REPORT

International Application No.

PC1/US 00/24373

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL 'Online!  ID: AV050323; Accession number: AV050323,  16 June 1999 (1999-06-16)  CARNINCI P. ET AL.: "Mus musculus adult  male pancreas cDNA, partial sequence,  clone 1810013H18"  XP002165897  abstract</p> <p style="text-align: center;">---</p>	1-5,7,11
A	<p>GALZIE Z. ET AL: "Fibroblast Growth  Factors and their receptors"  BIOCHEMISTRY AND CELL BIOLOGY,  vol. 75, no. 6, 1997, pages 669-685,  XP000925802  ISSN: 0829-8211</p> <p style="text-align: center;">---</p>	
A	<p>SMALLWOOD P.M. ET AL.: "Fibroblast Growth  Factor (FGF) homologous factors: new  members of the FGF family implicated in  nervous system development"  PROCEEDINGS OF THE NATIONAL ACADEMY OF  SCIENCES OF THE USA,  vol. 93, September 1996 (1996-09), pages  9850-9857, XP002058996  ISSN: 0027-8424</p> <p style="text-align: center;">-----</p>	

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 32-34 all in part

Present claims 32-34 (claims 33 and 34 in part) relate to a method involving an agonist/antagonist, without giving a true technical characterization of said agonist/antagonist.

The claims cover all agents having this agonist/antagonist effect, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such agents.

In fact, on page 53, present application recites that " an FGF-like polypeptide agonist or antagonist may be a protein, peptide, carbohydrate, lipid or small molecular weight molecule that interacts with FGF-like polypeptide to regulate its activity ". Moreover, on the same page 53, present application refers to antisense nucleic acid molecules as potential regulator of expression of the gene encoding the FGF-like polypeptide of the invention.

However, no specific carbohydrate, lipid or small molecular weight molecules are disclosed or defined in present application as having said agonist/antagonist effect. Moreover, concerning proteins and peptides, only antibodies against the polypeptide(s) disclosed can be derived from present application.

Therefore, in the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible.

Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define an agent by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible.

Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to antibodies raised against the polypeptide(s) of the invention and antisense molecules.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC1/US 00/24373

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0132678	A	10-05-2001	WO 0132678 A1	10-05-2001
			US 2001012628 A1	09-08-2001
WO 0138357	A	31-05-2001	AU 1628101 A	04-06-2001
			WO 0138357 A2	31-05-2001
WO 0136640	A	25-05-2001	AU 1922101 A	30-05-2001
			WO 0136640 A2	25-05-2001
WO 0118209	A	15-03-2001	AU 7368100 A	10-04-2001
			WO 0118209 A1	15-03-2001
WO 0149849	A	12-07-2001	AU 2631001 A	16-07-2001
			WO 0149849 A1	12-07-2001
WO 0054813	A	21-09-2000	AU 3755900 A	04-10-2000
			WO 0054813 A2	21-09-2000